

Software Engineering using Formal Methods

First-Order Logic

Wolfgang Ahrendt

26th September 2013

Install the KeY-Tool...

KeY used in Friday's exercise

Requires: Java \geq 5

Follow instructions on course page, under:

⇒ [Links, Papers, and Software](#)

We recommend using [Java Web Start](#):

- ▶ Start KeY with two clicks
(you need to trust our self-signed certificate)
- ▶ Java Web Start installed with standard JDK/JRE
- ▶ Usually browsers know filetype.
Otherwise open KeY.jnlp with javaws.

Install the KeY-Tool...

KeY used in Friday's exercise

Requires: Java \geq 5

Follow instructions on course page, under:

⇒ [Links, Papers, and Software](#)

We recommend using [Java Web Start](#):

- ▶ Start KeY with two clicks
(you need to trust our self-signed certificate)
- ▶ Java Web Start installed with standard JDK/JRE
- ▶ Usually browsers know filetype.
Otherwise open KeY.jnlp with javaws.

If you want to install KeY locally instead, download from www.key-project.org. For this course, install version 1.6.x.

Motivation for Introducing First-Order Logic

1) we specify JAVA programs with **Java Modeling Language (JML)**

JML combines

- ▶ JAVA expressions
- ▶ **First-Order Logic (FOL)**

Motivation for Introducing First-Order Logic

1) we specify JAVA programs with **Java Modeling Language (JML)**

JML combines

- ▶ JAVA expressions
- ▶ **First-Order Logic (FOL)**

2) we verify JAVA programs using **Dynamic Logic**

Dynamic Logic combines

- ▶ **First-Order Logic (FOL)**
- ▶ JAVA programs

we introduce:

- ▶ FOL as a language
- ▶ calculus for proving FOL formulas
- ▶ KeY system as propositional, and first-order, prover (for now)
- ▶ (formal semantics: if time)

First-Order Logic: Signature

Signature

A first-order signature Σ consists of

- ▶ a set T_Σ of types
- ▶ a set F_Σ of function symbols
- ▶ a set P_Σ of predicate symbols
- ▶ a typing α_Σ

First-Order Logic: Signature

Signature

A first-order signature Σ consists of

- ▶ a set T_Σ of types
- ▶ a set F_Σ of function symbols
- ▶ a set P_Σ of predicate symbols
- ▶ a typing α_Σ

intuitively, the typing α_Σ determines

- ▶ for each function and predicate symbol:
 - ▶ its arity, i.e., number of arguments
 - ▶ its argument types
- ▶ for each function symbol its result type.

First-Order Logic: Signature

Signature

A first-order signature Σ consists of

- ▶ a set T_Σ of types
- ▶ a set F_Σ of function symbols
- ▶ a set P_Σ of predicate symbols
- ▶ a typing α_Σ

intuitively, the typing α_Σ determines

- ▶ for each function and predicate symbol:
 - ▶ its arity, i.e., number of arguments
 - ▶ its argument types
- ▶ for each function symbol its result type.

formally:

- ▶ $\alpha_\Sigma(p) \in T_\Sigma^*$ for all $p \in P_\Sigma$ (arity of p is $|\alpha_\Sigma(p)|$)
- ▶ $\alpha_\Sigma(f) \in T_\Sigma^* \times T_\Sigma$ for all $f \in F_\Sigma$ (arity of f is $|\alpha_\Sigma(f)| - 1$)

Example Signature 1 + Constants

$$T_{\Sigma_1} = \{\text{int}\},$$

$$F_{\Sigma_1} = \{+, -\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\},$$

$$P_{\Sigma_1} = \{<\}$$

Example Signature 1 + Constants

$$T_{\Sigma_1} = \{\text{int}\},$$

$$F_{\Sigma_1} = \{+, -\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\},$$

$$P_{\Sigma_1} = \{<\}$$

$$\alpha_{\Sigma_1}(<) = (\text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(+) = \alpha_{\Sigma_1}(-) = (\text{int}, \text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(0) = \alpha_{\Sigma_1}(1) = \alpha_{\Sigma_1}(-1) = \dots = (\text{int})$$

Example Signature 1 + Constants

$$T_{\Sigma_1} = \{\text{int}\},$$

$$F_{\Sigma_1} = \{+, -\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\},$$

$$P_{\Sigma_1} = \{<\}$$

$$\alpha_{\Sigma_1}(<) = (\text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(+) = \alpha_{\Sigma_1}(-) = (\text{int}, \text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(0) = \alpha_{\Sigma_1}(1) = \alpha_{\Sigma_1}(-1) = \dots = (\text{int})$$

Constant Symbols

A function symbol f with $|\alpha_{\Sigma_1}(f)| = 1$ (i.e., with arity 0) is called *constant symbol*.

Example Signature 1 + Constants

$$T_{\Sigma_1} = \{\text{int}\},$$

$$F_{\Sigma_1} = \{+, -\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\},$$

$$P_{\Sigma_1} = \{<\}$$

$$\alpha_{\Sigma_1}(<) = (\text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(+) = \alpha_{\Sigma_1}(-) = (\text{int}, \text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(0) = \alpha_{\Sigma_1}(1) = \alpha_{\Sigma_1}(-1) = \dots = (\text{int})$$

Constant Symbols

A function symbol f with $|\alpha_{\Sigma_1}(f)| = 1$ (i.e., with arity 0) is called *constant symbol*.

here, the constant symbols are: $\dots, -2, -1, 0, 1, 2, \dots$

Type declaration of signature symbols

- ▶ Write τx ; to declare variable x of type τ
- ▶ Write $p(\tau_1, \dots, \tau_r)$; for $\alpha(p) = (\tau_1, \dots, \tau_r)$
- ▶ Write $\tau f(\tau_1, \dots, \tau_r)$; for $\alpha(f) = (\tau_1, \dots, \tau_r, \tau)$

$r = 0$ is allowed, then write f instead of $f()$, etc.

Syntax of First-Order Logic: Signature Cont'd

Type declaration of signature symbols

- ▶ Write τx ; to declare variable x of type τ
- ▶ Write $p(\tau_1, \dots, \tau_r)$; for $\alpha(p) = (\tau_1, \dots, \tau_r)$
- ▶ Write $\tau f(\tau_1, \dots, \tau_r)$; for $\alpha(f) = (\tau_1, \dots, \tau_r, \tau)$

$r = 0$ is allowed, then write f instead of $f()$, etc.

Example

Variables `integerArray a; int i;`

Predicate Symbols `isEmpty(List); alertOn;`

Function Symbols `int arrayLookup(int); Object o;`

Example Signature 1 + Notation

typing of Signature 1:

$$\alpha_{\Sigma_1}(<) = (\text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(+) = \alpha_{\Sigma_1}(-) = (\text{int}, \text{int}, \text{int})$$

$$\alpha_{\Sigma_1}(0) = \alpha_{\Sigma_1}(1) = \alpha_{\Sigma_1}(-1) = \dots = (\text{int})$$

can alternatively be written as:

```
<(int, int);
```

```
int +(int, int);
```

```
int 0; int 1; int -1; ...
```


Example Signature 2

$$T_{\Sigma_2} = \{\text{int}, \text{LinkedList}\},$$

$$F_{\Sigma_2} = \{\text{null}, \text{new}, \text{elem}, \text{next}\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\}$$

$$P_{\Sigma_2} = \{\}$$

Example Signature 2

$$T_{\Sigma_2} = \{\text{int}, \text{LinkedList}\},$$

$$F_{\Sigma_2} = \{\text{null}, \text{new}, \text{elem}, \text{next}\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\}$$

$$P_{\Sigma_2} = \{\}$$

intuitively, elem and next model fields of LinkedList objects

Example Signature 2

$$T_{\Sigma_2} = \{\text{int}, \text{LinkedList}\},$$

$$F_{\Sigma_2} = \{\text{null}, \text{new}, \text{elem}, \text{next}\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\}$$

$$P_{\Sigma_2} = \{\}$$

intuitively, elem and next model fields of LinkedList objects

type declarations:

Example Signature 2

$$T_{\Sigma_2} = \{\text{int}, \text{LinkedList}\},$$

$$F_{\Sigma_2} = \{\text{null}, \text{new}, \text{elem}, \text{next}\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\}$$

$$P_{\Sigma_2} = \{\}$$

intuitively, elem and next model fields of LinkedList objects

type declarations:

```
LinkedList null;  
LinkedList new(int,LinkedList);  
int elem(LinkedList);  
LinkedList next(LinkedList);
```

Example Signature 2

$$\begin{aligned}T_{\Sigma_2} &= \{\text{int}, \text{LinkedList}\}, \\F_{\Sigma_2} &= \{\text{null}, \text{new}, \text{elem}, \text{next}\} \cup \{\dots, -2, -1, 0, 1, 2, \dots\} \\P_{\Sigma_2} &= \{\}\end{aligned}$$

intuitively, elem and next model fields of LinkedList objects

type declarations:

```
LinkedList null;  
LinkedList new(int,LinkedList);  
int elem(LinkedList);  
LinkedList next(LinkedList);
```

and as before:

```
int 0; int 1; int -1; ...
```

First-Order Terms

We assume a set V of variables ($V \cap (F_\Sigma \cup P_\Sigma) = \emptyset$).
Each $v \in V$ has a unique type $\alpha_\Sigma(v) \in T_\Sigma$.

First-Order Terms

We assume a set V of variables ($V \cap (F_\Sigma \cup P_\Sigma) = \emptyset$).
Each $v \in V$ has a unique type $\alpha_\Sigma(v) \in T_\Sigma$.

Terms are defined recursively:

Terms

A first-order term of type $\tau \in T_\Sigma$

- ▶ is either a variable of type τ , or
- ▶ has the form $f(t_1, \dots, t_n)$,
where $f \in F_\Sigma$ has result type τ , and each t_i is term of the correct type, following the typing α_Σ of f .

First-Order Terms

We assume a set V of variables ($V \cap (F_\Sigma \cup P_\Sigma) = \emptyset$).
Each $v \in V$ has a unique type $\alpha_\Sigma(v) \in T_\Sigma$.

Terms are defined recursively:

Terms

A first-order term of type $\tau \in T_\Sigma$

- ▶ is either a variable of type τ , or
- ▶ has the form $f(t_1, \dots, t_n)$,
where $f \in F_\Sigma$ has result type τ , and each t_i is term of the correct type, following the typing α_Σ of f .

If f is a constant symbol, the term is written f , instead of $f()$.

Terms over Signature 1

example terms over Σ_1 :

(assume variables `int v1`; `int v2`;))

Terms over Signature 1

example terms over Σ_1 :

(assume variables $\text{int } v_1; \text{ int } v_2;$)

- ▶ -7
- ▶ $+(-2, 99)$
- ▶ $-(7, 8)$
- ▶ $+(-(7, 8), 1)$
- ▶ $+(-(v_1, 8), v_2)$

Terms over Signature 1

example terms over Σ_1 :

(assume variables $\text{int } v_1; \text{ int } v_2;$)

- ▶ -7
- ▶ $+(-2, 99)$
- ▶ $-(7, 8)$
- ▶ $+(-(7, 8), 1)$
- ▶ $+(-(v_1, 8), v_2)$

some variants of FOL allow infix notation of functions:

- ▶ $-2 + 99$
- ▶ $7 - 8$
- ▶ $(7 - 8) + 1$
- ▶ $(v_1 - 8) + v_2$

Terms over Signature 2

example terms over Σ_2 :

(assume variables `LinkedList` o ; `int` v ;))

Terms over Signature 2

example terms over Σ_2 :

(assume variables `LinkedList` o ; `int` v ;))

- ▶ `-7`
- ▶ `null`
- ▶ `new(13, null)`
- ▶ `elem(new(13, null))`
- ▶ `next(next(o))`

Terms over Signature 2

example terms over Σ_2 :

(assume variables `LinkedList` `o`; `int` `v`;

- ▶ `-7`
- ▶ `null`
- ▶ `new(13, null)`
- ▶ `elem(new(13, null))`
- ▶ `next(next(o))`

for first-order functions modeling object fields,
we allow dotted postfix notation:

- ▶ `new(13, null).elem`
- ▶ `o.next.next`

Atomic Formulas

Given a signature Σ .

An atomic formula has either of the forms

- ▶ *true*
- ▶ *false*
- ▶ $t_1 = t_2$ (“equality”),
where t_1 and t_2 are first-order terms of the same type.
- ▶ $p(t_1, \dots, t_n)$ (“predicate”),
where $p \in P_\Sigma$, and each t_i is term of the correct type,
following the typing α_Σ of p .

Atomic Formulas over Signature 1

example formulas over Σ_1 :
(assume variable `int v`;))

Atomic Formulas over Signature 1

example formulas over Σ_1 :
(assume variable `int v`;

- ▶ $7 = 8$
- ▶ $7 < 8$
- ▶ $-2 - v < 99$
- ▶ $v < (v + 1)$

Atomic Formulas over Signature 2

example formulas over Σ_2 :

(assume variables `LinkedList o; int v;`)

Atomic Formulas over Signature 2

example formulas over Σ_2 :

(assume variables `LinkedList o`; `int v`;))

- ▶ `new(13, null) = null`
- ▶ `elem(new(13, null)) = 13`
- ▶ `next(new(13, null)) = null`
- ▶ `next(next(o)) = o`

First-order Formulas

Formulas

- ▶ each atomic formula is a formula
- ▶ with ϕ and ψ formulas, x a variable, and τ a type, the following are also formulas:
 - ▶ $\neg\phi$ (“not ϕ ”)
 - ▶ $\phi \wedge \psi$ (“ ϕ and ψ ”)
 - ▶ $\phi \vee \psi$ (“ ϕ or ψ ”)
 - ▶ $\phi \rightarrow \psi$ (“ ϕ implies ψ ”)
 - ▶ $\phi \leftrightarrow \psi$ (“ ϕ is equivalent to ψ ”)

First-order Formulas

Formulas

- ▶ each atomic formula is a formula
- ▶ with ϕ and ψ formulas, x a variable, and τ a type, the following are also formulas:
 - ▶ $\neg\phi$ (“not ϕ ”)
 - ▶ $\phi \wedge \psi$ (“ ϕ and ψ ”)
 - ▶ $\phi \vee \psi$ (“ ϕ or ψ ”)
 - ▶ $\phi \rightarrow \psi$ (“ ϕ implies ψ ”)
 - ▶ $\phi \leftrightarrow \psi$ (“ ϕ is equivalent to ψ ”)
 - ▶ $\forall \tau x; \phi$ (“for all x of type τ holds ϕ ”)

First-order Formulas

Formulas

- ▶ each atomic formula is a formula
- ▶ with ϕ and ψ formulas, x a variable, and τ a type, the following are also formulas:
 - ▶ $\neg\phi$ (“not ϕ ”)
 - ▶ $\phi \wedge \psi$ (“ ϕ and ψ ”)
 - ▶ $\phi \vee \psi$ (“ ϕ or ψ ”)
 - ▶ $\phi \rightarrow \psi$ (“ ϕ implies ψ ”)
 - ▶ $\phi \leftrightarrow \psi$ (“ ϕ is equivalent to ψ ”)
 - ▶ $\forall \tau x; \phi$ (“for all x of type τ holds ϕ ”)
 - ▶ $\exists \tau x; \phi$ (“there exists an x of type τ such that ϕ ”)

First-order Formulas

Formulas

- ▶ each atomic formula is a formula
- ▶ with ϕ and ψ formulas, x a variable, and τ a type, the following are also formulas:
 - ▶ $\neg\phi$ (“not ϕ ”)
 - ▶ $\phi \wedge \psi$ (“ ϕ and ψ ”)
 - ▶ $\phi \vee \psi$ (“ ϕ or ψ ”)
 - ▶ $\phi \rightarrow \psi$ (“ ϕ implies ψ ”)
 - ▶ $\phi \leftrightarrow \psi$ (“ ϕ is equivalent to ψ ”)
 - ▶ $\forall \tau x; \phi$ (“for all x of type τ holds ϕ ”)
 - ▶ $\exists \tau x; \phi$ (“there exists an x of type τ such that ϕ ”)

In $\forall \tau x; \phi$ and $\exists \tau x; \phi$ the variable x is ‘bound’ (i.e., ‘not free’).

First-order Formulas

Formulas

- ▶ each atomic formula is a formula
- ▶ with ϕ and ψ formulas, x a variable, and τ a type, the following are also formulas:
 - ▶ $\neg\phi$ (“not ϕ ”)
 - ▶ $\phi \wedge \psi$ (“ ϕ and ψ ”)
 - ▶ $\phi \vee \psi$ (“ ϕ or ψ ”)
 - ▶ $\phi \rightarrow \psi$ (“ ϕ implies ψ ”)
 - ▶ $\phi \leftrightarrow \psi$ (“ ϕ is equivalent to ψ ”)
 - ▶ $\forall \tau x; \phi$ (“for all x of type τ holds ϕ ”)
 - ▶ $\exists \tau x; \phi$ (“there exists an x of type τ such that ϕ ”)

In $\forall \tau x; \phi$ and $\exists \tau x; \phi$ the variable x is ‘bound’ (i.e., ‘not free’).
Formulas with no free variable are ‘closed’.

First-order Formulas: Examples

(signatures/types left out here)

Example (There are at least two elements)



First-order Formulas: Examples

(signatures/types left out here)

Example (There are at least two elements)

$$\exists x, y; \neg(x = y)$$

First-order Formulas: Examples

(signatures/types left out here)

Example (Strict partial order)



First-order Formulas: Examples

(signatures/types left out here)

Example (Strict partial order)

Irreflexivity $\forall x; \neg(x < x)$

Asymmetry $\forall x; \forall y; (x < y \rightarrow \neg(y < x))$

Transitivity $\forall x; \forall y; \forall z;$
 $(x < y \wedge y < z \rightarrow x < z)$

First-order Formulas: Examples

(signatures/types left out here)

Example (Strict partial order)

Irreflexivity $\forall x; \neg(x < x)$

Asymmetry $\forall x; \forall y; (x < y \rightarrow \neg(y < x))$

Transitivity $\forall x; \forall y; \forall z;$
 $(x < y \wedge y < z \rightarrow x < z)$

(is any of the three formulas redundant?)

Semantics (briefly here, more thorough later)

Domain

A domain \mathcal{D} is a set of elements which are (potentially) the *meaning* of terms and variables.

Semantics (briefly here, more thorough later)

Domain

A domain \mathcal{D} is a set of elements which are (potentially) the *meaning* of terms and variables.

Interpretation

An interpretation \mathcal{I} (over \mathcal{D}) assigns *meaning* to the symbols in $F_\Sigma \cup P_\Sigma$ (assigning functions to function symbols, relations to predicate symbols).

Semantics (briefly here, more thorough later)

Domain

A domain \mathcal{D} is a set of elements which are (potentially) the *meaning* of terms and variables.

Interpretation

An interpretation \mathcal{I} (over \mathcal{D}) assigns *meaning* to the symbols in $F_\Sigma \cup P_\Sigma$ (assigning functions to function symbols, relations to predicate symbols).

Valuation

In a given \mathcal{D} and \mathcal{I} , a closed formula evaluates to either T or F .

Semantics (briefly here, more thorough later)

Domain

A domain \mathcal{D} is a set of elements which are (potentially) the *meaning* of terms and variables.

Interpretation

An interpretation \mathcal{I} (over \mathcal{D}) assigns *meaning* to the symbols in $F_\Sigma \cup P_\Sigma$ (assigning functions to function symbols, relations to predicate symbols).

Valuation

In a given \mathcal{D} and \mathcal{I} , a closed formula evaluates to either T or F .

Validity

A closed formula is **valid** if it evaluates to T in **all** \mathcal{D} and \mathcal{I} .

Semantics (briefly here, more thorough later)

Domain

A domain \mathcal{D} is a set of elements which are (potentially) the *meaning* of terms and variables.

Interpretation

An interpretation \mathcal{I} (over \mathcal{D}) assigns *meaning* to the symbols in $F_\Sigma \cup P_\Sigma$ (assigning functions to function symbols, relations to predicate symbols).

Valuation

In a given \mathcal{D} and \mathcal{I} , a closed formula evaluates to either T or F .

Validity

A closed formula is **valid** if it evaluates to T in **all** \mathcal{D} and \mathcal{I} .

In the context of specification/verification of programs:
each $(\mathcal{D}, \mathcal{I})$ is called a **'state'**.

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$

▶ $\neg(\phi \vee \psi) \leftrightarrow$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$

▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$
- ▶ $\neg(\text{false} \wedge \phi)$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$

▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$

▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$

▶ $(\text{false} \vee \phi) \leftrightarrow \phi$

▶ $\text{true} \vee \phi$

▶ $\neg(\text{false} \wedge \phi)$

▶ $(\phi \rightarrow \psi) \leftrightarrow$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$
- ▶ $\neg(\text{false} \wedge \phi)$
- ▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$
- ▶ $\neg(\text{false} \wedge \phi)$
- ▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$
- ▶ $\phi \rightarrow \text{true}$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$
- ▶ $\neg(\text{false} \wedge \phi)$
- ▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$
- ▶ $\phi \rightarrow \text{true}$
- ▶ $\text{false} \rightarrow \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$
- ▶ $\neg(\text{false} \wedge \phi)$
- ▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$
- ▶ $\phi \rightarrow \text{true}$
- ▶ $\text{false} \rightarrow \phi$
- ▶ $(\text{true} \rightarrow \phi) \leftrightarrow \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

- ▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$
- ▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$
- ▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$
- ▶ $(\text{false} \vee \phi) \leftrightarrow \phi$
- ▶ $\text{true} \vee \phi$
- ▶ $\neg(\text{false} \wedge \phi)$
- ▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$
- ▶ $\phi \rightarrow \text{true}$
- ▶ $\text{false} \rightarrow \phi$
- ▶ $(\text{true} \rightarrow \phi) \leftrightarrow \phi$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$

▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$

▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$

▶ $(\text{false} \vee \phi) \leftrightarrow \phi$

▶ $\text{true} \vee \phi$

▶ $\neg(\text{false} \wedge \phi)$

▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$

▶ $\phi \rightarrow \text{true}$

▶ $\text{false} \rightarrow \phi$

▶ $(\text{true} \rightarrow \phi) \leftrightarrow \phi$

▶ $(\phi \rightarrow \text{false}) \leftrightarrow$

Useful Valid Formulas

Let ϕ and ψ be arbitrary, closed formulas (whether valid or not).

The following formulas are valid:

▶ $\neg(\phi \wedge \psi) \leftrightarrow \neg\phi \vee \neg\psi$

▶ $\neg(\phi \vee \psi) \leftrightarrow \neg\phi \wedge \neg\psi$

▶ $(\text{true} \wedge \phi) \leftrightarrow \phi$

▶ $(\text{false} \vee \phi) \leftrightarrow \phi$

▶ $\text{true} \vee \phi$

▶ $\neg(\text{false} \wedge \phi)$

▶ $(\phi \rightarrow \psi) \leftrightarrow (\neg\phi \vee \psi)$

▶ $\phi \rightarrow \text{true}$

▶ $\text{false} \rightarrow \phi$

▶ $(\text{true} \rightarrow \phi) \leftrightarrow \phi$

▶ $(\phi \rightarrow \text{false}) \leftrightarrow \neg\phi$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists x \phi) \leftrightarrow \forall x \neg \phi$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists x; \phi) \leftrightarrow \forall x; \neg\phi$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow (\forall \tau x; \phi) \wedge (\forall \tau x; \psi)$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow (\forall \tau x; \phi) \wedge (\forall \tau x; \psi)$
- ▶ $(\exists \tau x; \phi \vee \psi) \leftrightarrow$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow (\forall \tau x; \phi) \wedge (\forall \tau x; \psi)$
- ▶ $(\exists \tau x; \phi \vee \psi) \leftrightarrow (\exists \tau x; \phi) \vee (\exists \tau x; \psi)$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow (\forall \tau x; \phi) \wedge (\forall \tau x; \psi)$
- ▶ $(\exists \tau x; \phi \vee \psi) \leftrightarrow (\exists \tau x; \phi) \vee (\exists \tau x; \psi)$

Are the following formulas also valid?

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow (\forall \tau x; \phi) \wedge (\forall \tau x; \psi)$
- ▶ $(\exists \tau x; \phi \vee \psi) \leftrightarrow (\exists \tau x; \phi) \vee (\exists \tau x; \psi)$

Are the following formulas also valid?

- ▶ $(\forall \tau x; \phi \vee \psi) \leftrightarrow (\forall \tau x; \phi) \vee (\forall \tau x; \psi)$

Useful Valid Formulas

Assume that x is the only variable which may appear freely in ϕ or ψ .

The following formulas are valid:

- ▶ $\neg(\exists \tau x; \phi) \leftrightarrow \forall \tau x; \neg\phi$
- ▶ $\neg(\forall \tau x; \phi) \leftrightarrow \exists \tau x; \neg\phi$
- ▶ $(\forall \tau x; \phi \wedge \psi) \leftrightarrow (\forall \tau x; \phi) \wedge (\forall \tau x; \psi)$
- ▶ $(\exists \tau x; \phi \vee \psi) \leftrightarrow (\exists \tau x; \phi) \vee (\exists \tau x; \psi)$

Are the following formulas also valid?

- ▶ $(\forall \tau x; \phi \vee \psi) \leftrightarrow (\forall \tau x; \phi) \vee (\forall \tau x; \psi)$
- ▶ $(\exists \tau x; \phi \wedge \psi) \leftrightarrow (\exists \tau x; \phi) \wedge (\exists \tau x; \psi)$

Remark on Concrete Syntax

	Text book	SPIN	KeY
Negation	\neg	!	!
Conjunction	\wedge	&&	&
Disjunction	\vee		
Implication	\rightarrow, \supset	\rightarrow	\rightarrow
Equivalence	\leftrightarrow	\leftrightarrow	\leftrightarrow
Universal Quantifier	$\forall x; \phi$	n/a	<code>\forall x; \phi</code>
Existential Quantifier	$\exists x; \phi$	n/a	<code>\exists x; \phi</code>
Value equality	=	==	=

Part I

Sequent Calculus for FOL

Reasoning by Syntactic Transformation

Prove Validity of ϕ by syntactic transformation of ϕ

Reasoning by Syntactic Transformation

Prove Validity of ϕ by **syntactic** transformation of ϕ

Logic Calculus: **Sequent Calculus** based on notion of **sequent**:

$$\underbrace{\psi_1, \dots, \psi_m}_{\text{Antecedent}} \Rightarrow \underbrace{\phi_1, \dots, \phi_n}_{\text{Succedent}}$$

Reasoning by Syntactic Transformation

Prove Validity of ϕ by syntactic transformation of ϕ

Logic Calculus: **Sequent Calculus** based on notion of **sequent**:

$$\underbrace{\psi_1, \dots, \psi_m}_{\text{Antecedent}} \Rightarrow \underbrace{\phi_1, \dots, \phi_n}_{\text{Succedent}}$$

has same meaning as

$$(\psi_1 \wedge \dots \wedge \psi_m) \rightarrow (\phi_1 \vee \dots \vee \phi_n)$$

Reasoning by Syntactic Transformation

Prove Validity of ϕ by syntactic transformation of ϕ

Logic Calculus: **Sequent Calculus** based on notion of **sequent**:

$$\underbrace{\psi_1, \dots, \psi_m}_{\text{Antecedent}} \Rightarrow \underbrace{\phi_1, \dots, \phi_n}_{\text{Succedent}}$$

has same meaning as

$$(\psi_1 \wedge \dots \wedge \psi_m) \rightarrow (\phi_1 \vee \dots \vee \phi_n)$$

which has (for closed formulas ψ_i, ϕ_i) same meaning as

$$\{\psi_1, \dots, \psi_m\} \models \phi_1 \vee \dots \vee \phi_n$$

Notation for Sequents

$$\psi_1, \dots, \psi_m \Rightarrow \phi_1, \dots, \phi_n$$

Consider antecedent/succedent as sets of formulas, may be empty

Notation for Sequents

$$\psi_1, \dots, \psi_m \Rightarrow \phi_1, \dots, \phi_n$$

Consider antecedent/succedent as sets of formulas, may be empty

Schema Variables

ϕ, ψ, \dots match formulas, Γ, Δ, \dots match sets of formulas

Characterize infinitely many sequents with single schematic sequent, e.g.,

$$\Gamma \Rightarrow \phi \wedge \psi, \Delta$$

Matches any sequent with occurrence of conjunction in succedent

Call $\phi \wedge \psi$ **main formula** and Γ, Δ **side formulas** of sequent

Any sequent of the form $\Gamma, \phi \Rightarrow \phi, \Delta$ is logically valid: **axiom**

Sequent Calculus Rules

Write syntactic transformation schema for sequents that reflects semantics of connectives as closely as possible

$$\text{RuleName} \frac{\overbrace{\Gamma_1 \Rightarrow \Delta_1 \quad \dots \quad \Gamma_r \Rightarrow \Delta_r}^{\text{Premises}}}{\underbrace{\Gamma \Rightarrow \Delta}_{\text{Conclusion}}}$$

Sequent Calculus Rules

Write syntactic transformation schema for sequents that reflects semantics of connectives as closely as possible

$$\text{RuleName} \frac{\overbrace{\Gamma_1 \Rightarrow \Delta_1 \quad \dots \quad \Gamma_r \Rightarrow \Delta_r}^{\text{Premises}}}{\underbrace{\Gamma \Rightarrow \Delta}_{\text{Conclusion}}}$$

Meaning: For proving the Conclusion, it suffices to prove all Premises.

Sequent Calculus Rules

Write syntactic transformation schema for sequents that reflects semantics of connectives as closely as possible

$$\text{RuleName} \frac{\overbrace{\Gamma_1 \Rightarrow \Delta_1 \quad \dots \quad \Gamma_r \Rightarrow \Delta_r}^{\text{Premises}}}{\underbrace{\Gamma \Rightarrow \Delta}_{\text{Conclusion}}}$$

Meaning: For proving the Conclusion, it suffices to prove all Premises.

Example

$$\text{andRight} \frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$$

Sequent Calculus Rules

Write syntactic transformation schema for sequents that reflects semantics of connectives as closely as possible

$$\text{RuleName} \frac{\overbrace{\Gamma_1 \Rightarrow \Delta_1 \quad \dots \quad \Gamma_r \Rightarrow \Delta_r}^{\text{Premises}}}{\underbrace{\Gamma \Rightarrow \Delta}_{\text{Conclusion}}}$$

Meaning: For proving the Conclusion, it suffices to prove all Premises.

Example

$$\text{andRight} \frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$$

Admissible to have no premisses (iff conclusion is valid, e.g., axiom)

Sequent Calculus Rules

Write syntactic transformation schema for sequents that reflects semantics of connectives as closely as possible

$$\text{RuleName} \frac{\overbrace{\Gamma_1 \Rightarrow \Delta_1 \quad \cdots \quad \Gamma_r \Rightarrow \Delta_r}^{\text{Premises}}}{\underbrace{\Gamma \Rightarrow \Delta}_{\text{Conclusion}}}$$

Meaning: For proving the Conclusion, it suffices to prove all Premises.

Example

$$\text{andRight} \frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$$

Admissible to have no premisses (iff conclusion is valid, e.g., axiom)

A rule is **sound** (correct) iff the validity of its premisses implies the validity of its conclusion.

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$
or	$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta}$

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$
or	$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta}$
imp	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \rightarrow \psi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta}$

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$
or	$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta}$
imp	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \rightarrow \psi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta}$
close	$\frac{}{\Gamma, \phi \Rightarrow \phi, \Delta}$	

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$
or	$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta}$
imp	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \rightarrow \psi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta}$
close	$\frac{}{\Gamma, \phi \Rightarrow \phi, \Delta}$	true $\frac{}{\Gamma \Rightarrow \text{true}, \Delta}$

'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$
or	$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta}$
imp	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \rightarrow \psi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta}$
close	$\frac{}{\Gamma, \phi \Rightarrow \phi, \Delta}$	true $\frac{}{\Gamma \Rightarrow \text{true}, \Delta}$ false $\frac{}{\Gamma, \text{false} \Rightarrow \Delta}$

Sequent Calculus Proofs

Goal to prove: $\mathcal{G} = \psi_1, \dots, \psi_m \Rightarrow \phi_1, \dots, \phi_n$

- ▶ find rule \mathcal{R} whose conclusion **matches** \mathcal{G}
- ▶ instantiate \mathcal{R} such that its conclusion is **identical** to \mathcal{G}
- ▶ apply that instantiation to all premisses of \mathcal{R} , resulting in new goals $\mathcal{G}_1, \dots, \mathcal{G}_r$
- ▶ recursively find proofs for $\mathcal{G}_1, \dots, \mathcal{G}_r$
- ▶ tree structure with goal as root
- ▶ **close** proof branch when rule without premiss encountered


Sequent Calculus Proofs

Goal to prove: $\mathcal{G} = \psi_1, \dots, \psi_m \Rightarrow \phi_1, \dots, \phi_n$

- ▶ find rule \mathcal{R} whose conclusion **matches** \mathcal{G}
- ▶ instantiate \mathcal{R} such that its conclusion is **identical** to \mathcal{G}
- ▶ apply that instantiation to all premisses of \mathcal{R} , resulting in new goals $\mathcal{G}_1, \dots, \mathcal{G}_r$
- ▶ recursively find proofs for $\mathcal{G}_1, \dots, \mathcal{G}_r$
- ▶ tree structure with goal as root
- ▶ **close** proof branch when rule without premiss encountered

Goal-directed proof search

In KeY tool proof displayed as JAVA Swing tree



A Simple Proof

$$\frac{\begin{array}{c} \text{---} \quad \text{---} \\ \hline \hline \hline \end{array}}{\Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q}$$

A Simple Proof

$$\frac{\frac{\frac{}{p \wedge (p \rightarrow q)} \Rightarrow q}{\Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q}}{\Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q}}$$

A Simple Proof

$$\frac{\frac{\frac{}{p, (p \rightarrow q) \Rightarrow q}}{p \wedge (p \rightarrow q) \Rightarrow q}}{\Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q}}$$

A Simple Proof

$$\frac{\frac{\frac{}{p \Rightarrow p, q}}{p, (p \rightarrow q) \Rightarrow q}}{p \wedge (p \rightarrow q) \Rightarrow q}}{\Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q}$$

A Simple Proof

$$\begin{array}{c} \text{CLOSE} \frac{*}{p \Rightarrow p, q} \quad \frac{*}{p, q \Rightarrow q} \text{CLOSE} \\ \hline p, (p \rightarrow q) \Rightarrow q \\ \hline p \wedge (p \rightarrow q) \Rightarrow q \\ \hline \Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q \end{array}$$

A Simple Proof

$$\frac{\frac{\text{CLOSE} \frac{*}{p \Rightarrow p, q}}{p, (p \rightarrow q) \Rightarrow q} \quad \frac{*}{p, q \Rightarrow q} \text{CLOSE}}{p \wedge (p \rightarrow q) \Rightarrow q}}{\Rightarrow (p \wedge (p \rightarrow q)) \rightarrow q}$$

A proof is **closed** iff all its branches are closed

Demo

prop.key

Proving Validity of First-Order Formulas

Proving a universally quantified formula

Claim: $\forall x; \phi$ is true

How is such a claim proved in mathematics?

Proving Validity of First-Order Formulas

Proving a universally quantified formula

Claim: $\forall x; \phi$ is true

How is such a claim proved in mathematics?

All even numbers are divisible by 2 $\forall \text{int } x; (\text{even}(x) \rightarrow \text{divByTwo}(x))$

Proving Validity of First-Order Formulas

Proving a universally quantified formula

Claim: $\forall x; \phi$ is true

How is such a claim proved in mathematics?

All even numbers are divisible by 2 $\forall \text{int } x; (\text{even}(x) \rightarrow \text{divByTwo}(x))$

Let c be an arbitrary number Declare “unused” constant `int c`

Proving Validity of First-Order Formulas

Proving a universally quantified formula

Claim: $\forall x; \phi$ is true

How is such a claim proved in mathematics?

All even numbers are divisible by 2 $\forall \text{int } x; (\text{even}(x) \rightarrow \text{divByTwo}(x))$

Let c be an arbitrary number Declare “unused” constant `int c`

The even number c is divisible by 2 prove `even(c) \rightarrow divByTwo(c)`

Proving Validity of First-Order Formulas

Proving a universally quantified formula

Claim: $\forall \tau x; \phi$ is true

How is such a claim proved in mathematics?

All even numbers are divisible by 2 $\forall \text{int } x; (\text{even}(x) \rightarrow \text{divByTwo}(x))$

Let c be an arbitrary number Declare “unused” constant `int c`

The even number c is divisible by 2 prove `even(c) \rightarrow divByTwo(c)`

Sequent rule \forall -right

$$\text{forallRight} \frac{\Gamma \Rightarrow [x/c] \phi, \Delta}{\Gamma \Rightarrow \forall \tau x; \phi, \Delta}$$

- ▶ $[x/c] \phi$ is result of replacing each occurrence of x in ϕ with c
- ▶ c **new** constant of type τ

Proving Validity of First-Order Formulas Cont'd

Proving an existentially quantified formula

Claim: $\exists x; \phi$ is true

How is such a claim proved in mathematics?

Proving Validity of First-Order Formulas Cont'd

Proving an existentially quantified formula

Claim: $\exists \tau x; \phi$ is true

How is such a claim proved in mathematics?

There is at least one prime number $\exists \text{int } x; \text{prime}(x)$

Proving Validity of First-Order Formulas Cont'd

Proving an existentially quantified formula

Claim: $\exists \tau x; \phi$ is true

How is such a claim proved in mathematics?

There is at least one prime number $\exists \text{int } x; \text{prime}(x)$

Provide any "witness", say, 7 Use variable-free term `int 7`

Proving Validity of First-Order Formulas Cont'd

Proving an existentially quantified formula

Claim: $\exists x; \phi$ is true

How is such a claim proved in mathematics?

There is at least one prime number $\exists \text{int } x; \text{prime}(x)$

Provide any "witness", say, 7 Use variable-free term `int 7`

7 is a prime number `prime(7)`

Proving Validity of First-Order Formulas Cont'd

Proving an existentially quantified formula

Claim: $\exists \tau x; \phi$ is true

How is such a claim proved in mathematics?

There is at least one prime number $\exists \text{int } x; \text{prime}(x)$

Provide any "witness", say, 7 Use variable-free term `int 7`

7 is a prime number `prime(7)`

Sequent rule \exists -right

$$\text{existsRight} \frac{\Gamma \Rightarrow [x/t] \phi, \exists \tau x; \phi, \Delta}{\Gamma \Rightarrow \exists \tau x; \phi, \Delta}$$

- ▶ t any variable-free term of type τ
- ▶ Proof might not work with t ! Need to keep premise to try again

Proving Validity of First-Order Formulas Cont'd

Using a universally quantified formula

We assume $\forall x; \phi$ is true

How is such a fact **used** in a mathematical proof?

Proving Validity of First-Order Formulas Cont'd

Using a universally quantified formula

We assume $\forall x; \phi$ is true

How is such a fact **used** in a mathematical proof?

We know that all primes are odd $\forall \text{int } x; (\text{prime}(x) \rightarrow \text{odd}(x))$

Proving Validity of First-Order Formulas Cont'd

Using a universally quantified formula

We assume $\forall x; \phi$ is true

How is such a fact **used** in a mathematical proof?

We know that all primes are odd $\forall \text{int } x; (\text{prime}(x) \rightarrow \text{odd}(x))$

In particular, this holds for 17 Use variable-free term `int 17`

Proving Validity of First-Order Formulas Cont'd

Using a universally quantified formula

We assume $\forall x; \phi$ is true

How is such a fact **used** in a mathematical proof?

We know that all primes are odd $\forall \text{int } x; (\text{prime}(x) \rightarrow \text{odd}(x))$

In particular, this holds for 17 Use variable-free term `int 17`

We know: if 17 is prime it is odd $\text{prime}(17) \rightarrow \text{odd}(17)$

Proving Validity of First-Order Formulas Cont'd

Using a universally quantified formula

We assume $\forall \tau x; \phi$ is true

How is such a fact **used** in a mathematical proof?

We know that all primes are odd $\forall \text{int } x; (\text{prime}(x) \rightarrow \text{odd}(x))$

In particular, this holds for 17 Use variable-free term `int 17`

We know: if 17 is prime it is odd $\text{prime}(17) \rightarrow \text{odd}(17)$

Sequent rule \forall -left

$$\text{forallLeft} \frac{\Gamma, \forall \tau x; \phi, [x/t'] \phi \Rightarrow \Delta}{\Gamma, \forall \tau x; \phi \Rightarrow \Delta}$$

- ▶ t' any variable-free term of type τ
- ▶ We might need other instances besides t' ! Keep premise $\forall \tau x; \phi$

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

We assume $\exists x; \phi$ is true

How is such a fact **used** in a mathematical proof?

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

We assume $\exists x \phi$ is true

How is such a fact **used** in a mathematical proof?

We know such an element exists. Let's give it a new name for future reference.

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

We assume $\exists \tau x; \phi$ is true

How is such a fact **used** in a mathematical proof?

We know such an element exists. Let's give it a new name for future reference.

Sequent rule \exists -left

$$\text{existsLeft} \frac{\Gamma, [x/c] \phi \Rightarrow \Delta}{\Gamma, \exists \tau x; \phi \Rightarrow \Delta}$$

- ▶ c **new** constant of type τ

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Let x, y denote integer constants, both are not zero.

$$\neg(x = 0), \neg(y = 0) \quad \Rightarrow$$

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Let x, y denote integer constants, both are not zero. We know further that x divides y .

$$\neg(x = 0), \neg(y = 0), \exists \text{int } k; k * x = y \implies$$

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Let x, y denote integer constants, both are not zero. We know further that x divides y .

Show: $(y/x) * x = y$ ('/' is division on integers, i.e. the equation is not always true, e.g. $x = 2, y = 1$)

$$\neg(x = 0), \neg(y = 0), \exists \text{int } k; k * x = y \implies (y/x) * x = y$$

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Let x, y denote integer constants, both are not zero. We know further that x divides y .

Show: $(y/x) * x = y$ ('/' is division on integers, i.e. the equation is not always true, e.g. $x = 2, y = 1$)

Proof: We know x divides y , i.e. there exists a k such that $k * x = y$.

Let now c denote such a k .

$$\frac{\neg(x = 0), \neg(y = 0), c * x = y \implies (y/x) * x = y}{\neg(x = 0), \neg(y = 0), \exists \text{int } k; k * x = y \implies (y/x) * x = y}$$

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Let x, y denote integer constants, both are not zero. We know further that x divides y .

Show: $(y/x) * x = y$ ('/' is division on integers, i.e. the equation is not always true, e.g. $x = 2, y = 1$)

Proof: We know x divides y , i.e. there exists a k such that $k * x = y$. Let now c denote such a k . Hence we can replace y by $c * x$ on the right side (see slide 35).

$$\frac{\frac{\neg(x = 0), \neg(y = 0), c * x = y \implies ((c * x)/x) * x = y}{\neg(x = 0), \neg(y = 0), c * x = y \implies (y/x) * x = y}}{\neg(x = 0), \neg(y = 0), \exists \text{int } k; k * x = y \implies (y/x) * x = y}$$

Proving Validity of First-Order Formulas Cont'd

Using an existentially quantified formula

Let x, y denote integer constants, both are not zero. We know further that x divides y .

Show: $(y/x) * x = y$ ('/' is division on integers, i.e. the equation is not always true, e.g. $x = 2, y = 1$)

Proof: We know x divides y , i.e. there exists a k such that $k * x = y$. Let now c denote such a k . Hence we can replace y by $c * x$ on the right side (see slide 35). ... \square

$$\begin{array}{c} * \\ \hline \vdots \\ \hline \frac{\neg(x = 0), \neg(y = 0), c * x = y \implies ((c * x)/x) * x = y}{\neg(x = 0), \neg(y = 0), c * x = y \implies (y/x) * x = y} \\ \hline \frac{\neg(x = 0), \neg(y = 0), \exists \text{int } k; k * x = y \implies (y/x) * x = y}{\neg(x = 0), \neg(y = 0), \exists \text{int } k; k * x = y \implies (y/x) * x = y} \end{array}$$

Example (A simple theorem about binary relations)

$$\exists x; \forall y; p(x, y) \implies \forall y; \exists x; p(x, y)$$

Untyped logic: let static type of x and y be \top

Proving Validity of First-Order Formulas Cont'd

Example (A simple theorem about binary relations)

$$\frac{\forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y)}{\exists x; \forall y; p(x, y) \Rightarrow \forall y; \exists x; p(x, y)}$$

\exists -left: substitute **new** constant c of type \top for x

Proving Validity of First-Order Formulas Cont'd

Example (A simple theorem about binary relations)

$$\frac{\frac{\forall y; p(c, y) \Rightarrow \exists x; p(x, d)}{\forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y)}}{\exists x; \forall y; p(x, y) \Rightarrow \forall y; \exists x; p(x, y)}$$

\forall -right: substitute **new** constant d of type \top for y

Proving Validity of First-Order Formulas Cont'd

Example (A simple theorem about binary relations)

$$\frac{\frac{\frac{p(c, d), \forall y; p(c, y) \Rightarrow \exists x; p(x, d)}{\forall y; p(c, y) \Rightarrow \exists x; p(x, d)}}{\forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y)}}{\exists x; \forall y; p(x, y) \Rightarrow \forall y; \exists x; p(x, y)}$$

\forall -left: free to substitute **any** term of type \top for y , choose d

Proving Validity of First-Order Formulas Cont'd

Example (A simple theorem about binary relations)

$$\frac{\frac{\frac{p(c, d), \forall y; p(c, y) \Rightarrow p(c, d), \exists x; p(x, y)}{p(c, d), \forall y; p(c, y) \Rightarrow \exists x; p(x, d)}}{\forall y; p(c, y) \Rightarrow \exists x; p(x, d)}}{\forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y)} \\ \frac{\forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y)}{\exists x; \forall y; p(x, y) \Rightarrow \forall y; \exists x; p(x, y)}$$

\exists -right: free to substitute **any** term of type \top for x , choose c

Proving Validity of First-Order Formulas Cont'd

Example (A simple theorem about binary relations)

$$\begin{array}{c} * \\ \hline p(c, d), \forall y; p(c, y) \Rightarrow p(c, d), \exists x; p(x, y) \\ \hline p(c, d), \forall y; p(c, y) \Rightarrow \exists x; p(x, d) \\ \hline \forall y; p(c, y) \Rightarrow \exists x; p(x, d) \\ \hline \forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y) \\ \hline \exists x; \forall y; p(x, y) \Rightarrow \forall y; \exists x; p(x, y) \end{array}$$

Close

Proving Validity of First-Order Formulas Cont'd

Example (A simple theorem about binary relations)

$$\begin{array}{c} * \\ \hline p(c, d), \forall y; p(c, y) \Rightarrow p(c, d), \exists x; p(x, y) \\ \hline p(c, d), \forall y; p(c, y) \Rightarrow \exists x; p(x, d) \\ \hline \forall y; p(c, y) \Rightarrow \exists x; p(x, d) \\ \hline \forall y; p(c, y) \Rightarrow \forall y; \exists x; p(x, y) \\ \hline \exists x; \forall y; p(x, y) \Rightarrow \forall y; \exists x; p(x, y) \end{array}$$

Demo

relSimple.key

Proving Validity of First-Order Formulas Cont'd

Using an equation between terms

We assume $t = t'$ is true

How is such a fact used in a mathematical proof?

Proving Validity of First-Order Formulas Cont'd

Using an equation between terms

We assume $t = t'$ is true

How is such a fact used in a mathematical proof?

Use $x = y - 1$ to simplify $x + 1/y$ $x = y - 1 \Rightarrow 1 = x + 1/y$

Proving Validity of First-Order Formulas Cont'd

Using an equation between terms

We assume $t = t'$ is true

How is such a fact used in a mathematical proof?

Use $x = y - 1$ to simplify $x + 1/y$ $x = y - 1 \Rightarrow 1 = x + 1/y$

Replace x in conclusion with right-hand side of equation

Proving Validity of First-Order Formulas Cont'd

Using an equation between terms

We assume $t = t'$ is true

How is such a fact used in a mathematical proof?

Use $x = y-1$ to simplify $x+1/y$ $x = y-1 \Rightarrow 1 = x+1/y$

Replace x in conclusion with right-hand side of equation

We know: $x+1/y$ equal to $y-1+1/y$ $x = y-1 \Rightarrow 1 = y-1+1/y$

Proving Validity of First-Order Formulas Cont'd

Using an equation between terms

We assume $t = t'$ is true

How is such a fact used in a mathematical proof?

Use $x = y-1$ to simplify $x+1/y$ $x = y-1 \Rightarrow 1 = x+1/y$

Replace x in conclusion with right-hand side of equation

We know: $x+1/y$ equal to $y-1+1/y$ $x = y-1 \Rightarrow 1 = y-1+1/y$

Sequent rule =-left

$$\text{applyEqL} \frac{\Gamma, t = t', [t/t'] \phi \Rightarrow \Delta}{\Gamma, t = t', \phi \Rightarrow \Delta} \quad \text{applyEqR} \frac{\Gamma, t = t' \Rightarrow [t/t'] \phi, \Delta}{\Gamma, t = t' \Rightarrow \phi, \Delta}$$

- ▶ Always replace left- with right-hand side (use **eqSymm** if necessary)
- ▶ t, t' variable-free terms of the same type

Proving Validity of First-Order Formulas Cont'd

Closing a subgoal in a proof

- ▶ We derived a sequent that is obviously valid

$$\text{close } \frac{}{\Gamma, \phi \Rightarrow \phi, \Delta} \quad \text{true } \frac{}{\Gamma \Rightarrow \text{true}, \Delta} \quad \text{false } \frac{}{\Gamma, \text{false} \Rightarrow \Delta}$$

- ▶ We derived an **equation** that is obviously valid

$$\text{eqClose } \frac{}{\Gamma \Rightarrow t = t, \Delta}$$

Sequent Calculus for FOL at One Glance

	left side, antecedent	right side, succedent
\forall	$\frac{\Gamma, \forall \tau x; \phi, [x/t'] \phi \Rightarrow \Delta}{\Gamma, \forall \tau x; \phi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow [x/c] \phi, \Delta}{\Gamma \Rightarrow \forall \tau x; \phi, \Delta}$
\exists	$\frac{\Gamma, [x/c] \phi \Rightarrow \Delta}{\Gamma, \exists \tau x; \phi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow [x/t'] \phi, \exists \tau x; \phi, \Delta}{\Gamma \Rightarrow \exists \tau x; \phi, \Delta}$
$=$	$\frac{\Gamma, t = t' \Rightarrow [t/t'] \phi, \Delta}{\Gamma, t = t' \Rightarrow \phi, \Delta}$	$\frac{}{\Gamma \Rightarrow t = t, \Delta}$
	(+ application rule on left side)	

- ▶ $[t/t'] \phi$ is result of replacing each occurrence of t in ϕ with t'
- ▶ t, t' variable-free terms of type τ
- ▶ c **new** constant of type τ (occurs not on current proof branch)
- ▶ Equations can be reversed by commutativity

Recap: 'Propositional' Sequent Calculus Rules

main	left side (antecedent)	right side (succedent)
not	$\frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma, \neg\phi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma \Rightarrow \neg\phi, \Delta}$
and	$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta}$
or	$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta}$	$\frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta}$
imp	$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \rightarrow \psi \Rightarrow \Delta}$	$\frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta}$
close	$\frac{}{\Gamma, \phi \Rightarrow \phi, \Delta}$	true $\frac{}{\Gamma \Rightarrow \text{true}, \Delta}$ false $\frac{}{\Gamma, \text{false} \Rightarrow \Delta}$

Features of the KeY Theorem Prover

Demo

`rel.key, twoInstances.key`

Feature List

- ▶ Can work on multiple proofs simultaneously (task list)
- ▶ Proof trees visualized as JAVA Swing tree
- ▶ Point-and-click navigation within proof
- ▶ Undo proof steps, prune proof trees
- ▶ Pop-up menu with proof rules applicable in pointer focus
- ▶ Preview of rule effect as tool tip
- ▶ Quantifier instantiation and equality rules by drag-and-drop
- ▶ Possible to hide (and unhide) parts of a sequent
- ▶ Saving and loading of proofs

Literature for this Lecture

essential:

- ▶ W. Ahrendt
Using KeY
Chapter 10 in [KeYbook]

further reading:

- ▶ M. Giese
First-Order Logic
Chapter 2 in [KeYbook]

KeYbook B. Beckert, R. Hähnle, and P. Schmitt, editors, **Verification of Object-Oriented Software: The KeY Approach**, vol 4334 of *LNCS* (Lecture Notes in Computer Science), Springer, 2006 (access via Chalmers library → E-books → Lecture Notes in Computer Science)

Part II

First-Order Semantics

First-Order Semantics

From propositional to first-order semantics

- ▶ In prop. logic, an interpretation of variables with $\{T, F\}$ sufficed
- ▶ In first-order logic we must assign meaning to:
 - ▶ variables bound in quantifiers
 - ▶ constant and function symbols
 - ▶ predicate symbols
- ▶ Each variable or function value may denote a different item
- ▶ Respect typing: `int i`, `List l` **must** denote different items

What we need (to interpret a first-order formula)

1. A collection of **typed universes** of items
2. A mapping from **variables** to items
3. A mapping from **function** arguments to function values
4. The set of argument tuples where a **predicate** is true

First-Order Domains/Universes

1. A collection of **typed universes** of items

Definition (Universe/Domain)

A non-empty set \mathcal{D} of items is a **universe** or **domain**

Each element of \mathcal{D} has a fixed type given by $\delta : \mathcal{D} \rightarrow \mathcal{T}$

- ▶ Notation for the domain elements of type $\tau \in \mathcal{T}$:

$$\mathcal{D}^\tau = \{d \in \mathcal{D} \mid \delta(d) = \tau\}$$

- ▶ Each type $\tau \in \mathcal{T}$ must 'contain' at least one domain element:

$$\mathcal{D}^\tau \neq \emptyset$$

First-Order States

3. A mapping from function arguments to function values
4. The set of argument tuples where a predicate is true

Definition (First-Order State)

Let \mathcal{D} be a domain with typing function δ

Let f be declared as $\tau f(\tau_1, \dots, \tau_r)$;

Let p be declared as $p(\tau_1, \dots, \tau_r)$;

Let $\mathcal{I}(f) : \mathcal{D}^{\tau_1} \times \dots \times \mathcal{D}^{\tau_r} \rightarrow \mathcal{D}^{\tau}$

Let $\mathcal{I}(p) \subseteq \mathcal{D}^{\tau_1} \times \dots \times \mathcal{D}^{\tau_r}$

Then $\mathcal{S} = (\mathcal{D}, \delta, \mathcal{I})$ is a **first-order state**

First-Order States Cont'd

Example

Signature: `int i; short j; int f(int); Object obj; <(int,int);`

$\mathcal{D} = \{17, 2, o\}$ where all numbers are short

$$\mathcal{I}(i) = 17$$

$$\mathcal{I}(j) = 17$$

$$\mathcal{I}(obj) = o$$

\mathcal{D}^{int}	$\mathcal{I}(f)$
2	2
17	2

$\mathcal{D}^{\text{int}} \times \mathcal{D}^{\text{int}}$	in $\mathcal{I}(<)$?
(2, 2)	<i>F</i>
(2, 17)	<i>T</i>
(17, 2)	<i>F</i>
(17, 17)	<i>F</i>

One of uncountably many possible first-order states!

Semantics of Reserved Signature Symbols

Definition

Equality symbol $=$ declared as $= (\top, \top)$

Interpretation is fixed as $\mathcal{I}(=) = \{(d, d) \mid d \in \mathcal{D}\}$

“Referential Equality” (holds if arguments refer to identical item)

Exercise: write down the predicate table for example domain

Signature Symbols vs. Domain Elements

- ▶ Domain elements different from the terms representing them
- ▶ First-order formulas and terms have **no access** to domain

Example

Signature: Object obj1, obj2;

Domain: $\mathcal{D} = \{o\}$

In this state, necessarily $\mathcal{I}(\text{obj1}) = \mathcal{I}(\text{obj2}) = o$

Variable Assignments

2. A mapping from variables to objects

Think of variable assignment as environment for storage of local variables

Definition (Variable Assignment)

A **variable assignment** β maps variables to domain elements

It respects the variable type, i.e., if x has type τ then $\beta(x) \in \mathcal{D}^\tau$

Definition (Modified Variable Assignment)

Let y be variable of type τ , β variable assignment, $d \in \mathcal{D}^\tau$:

$$\beta_y^d(x) := \begin{cases} \beta(x) & x \neq y \\ d & x = y \end{cases}$$

Semantic Evaluation of Terms

Given a first-order state \mathcal{S} and a variable assignment β it is possible to evaluate first-order terms under \mathcal{S} and β

Definition (Valuation of Terms)

$val_{\mathcal{S},\beta} : \text{Term} \rightarrow \mathcal{D}$ such that $val_{\mathcal{S},\beta}(t) \in \mathcal{D}^\tau$ for $t \in \text{Term}_\tau$:

- ▶ $val_{\mathcal{S},\beta}(x) = \beta(x)$
- ▶ $val_{\mathcal{S},\beta}(f(t_1, \dots, t_r)) = \mathcal{I}(f)(val_{\mathcal{S},\beta}(t_1), \dots, val_{\mathcal{S},\beta}(t_r))$

Semantic Evaluation of Terms Cont'd

Example

Signature: `int i; short j; int f(int);`

$\mathcal{D} = \{17, 2, o\}$ where all numbers are short

Variables: Object `obj`; `int x`;

$$I(i) = 17$$

$$I(j) = 17$$

\mathcal{D}^{int}	$I(f)$
2	17
17	2

Var	β
<code>obj</code>	<code>o</code>
<code>x</code>	17

- ▶ $val_{S,\beta}(f(f(i)))$?
- ▶ $val_{S,\beta}(x)$?

Semantic Evaluation of Formulas

Definition (Valuation of Formulas)

$val_{S,\beta}(\phi)$ for $\phi \in For$

- ▶ $val_{S,\beta}(p(t_1, \dots, t_r)) = T$ iff $(val_{S,\beta}(t_1), \dots, val_{S,\beta}(t_r)) \in \mathcal{I}(p)$
- ▶ $val_{S,\beta}(\phi \wedge \psi) = T$ iff $val_{S,\beta}(\phi) = T$ and $val_{S,\beta}(\psi) = T$
- ▶ ... as in propositional logic
- ▶ $val_{S,\beta}(\forall \tau x; \phi) = T$ iff $val_{S,\beta_x^d}(\forall \tau x; \phi) = T$ for all $d \in \mathcal{D}^\tau$
- ▶ $val_{S,\beta}(\exists \tau x; \phi) = T$ iff $val_{S,\beta_x^d}(\exists \tau x; \phi) = T$ for at least one $d \in \mathcal{D}^\tau$

Semantic Evaluation of Formulas Cont'd

Example

Signature: `short j`; `int f(int)`; `Object obj`; `<(int,int)`;

$\mathcal{D} = \{17, 2, o\}$ where all numbers are short

$$\begin{aligned} \mathcal{I}(j) &= 17 \\ \mathcal{I}(\text{obj}) &= o \end{aligned}$$

\mathcal{D}^{int}	$\mathcal{I}(f)$
2	2
17	2

$\mathcal{D}^{\text{int}} \times \mathcal{D}^{\text{int}}$	in $\mathcal{I}(<)$?
(2, 2)	F
(2, 17)	T
(17, 2)	F
(17, 17)	F

- ▶ $\text{val}_{\mathcal{S},\beta}(f(j) < j)$?
- ▶ $\text{val}_{\mathcal{S},\beta}(\exists \text{int } x; f(x) = x)$?
- ▶ $\text{val}_{\mathcal{S},\beta}(\forall \text{Object } o1; \forall \text{Object } o2; o1 = o2)$?

Semantic Notions

Definition (Satisfiability, Truth, Validity)

$val_{S,\beta}(\phi) = T$		(ϕ is satisfiable)
$\mathcal{S} \models \phi$	iff for all $\beta : val_{S,\beta}(\phi) = T$	(ϕ is true in \mathcal{S})
$\models \phi$	iff for all $\mathcal{S} : \mathcal{S} \models \phi$	(ϕ is valid)

Closed formulas that are satisfiable are also true: one top-level notion

Semantic Notions

Definition (Satisfiability, Truth, Validity)

$$\begin{array}{lll} \text{val}_{\mathcal{S},\beta}(\phi) = T & & (\phi \text{ is } \mathbf{satisfiable}) \\ \mathcal{S} \models \phi & \text{iff for all } \beta : \text{val}_{\mathcal{S},\beta}(\phi) = T & (\phi \text{ is } \mathbf{true} \text{ in } \mathcal{S}) \\ \models \phi & \text{iff for all } \mathcal{S} : \mathcal{S} \models \phi & (\phi \text{ is } \mathbf{valid}) \end{array}$$

Closed formulas that are satisfiable are also true: one top-level notion

Example

- ▶ $f(j) < j$ is true in \mathcal{S}
- ▶ $\exists \text{int } x; i = x$ is valid
- ▶ $\exists \text{int } x; \neg(x = x)$ is not satisfiable