

Homotopy canonicity for cubical type theory

Thierry Coquand

Department of Computer Science and Engineering, University of Gothenburg, Sweden
coquand@chalmers.se

Simon Huber

Department of Computer Science and Engineering, University of Gothenburg, Sweden
simon.huber@cse.gu.se

Christian Sattler

Department of Computer Science and Engineering, University of Gothenburg, Sweden
sattler@chalmers.se

Abstract

Cubical type theory provides a constructive justification of homotopy type theory and satisfies canonicity: every natural number is convertible to a numeral. A crucial ingredient of cubical type theory is a path lifting operation which is explained computationally by induction on the type involving several non-canonical choices. In this paper we show by a scoping argument that if we remove these equations for the path lifting operation from the system, we still retain *homotopy* canonicity: every natural number is *path equal* to a numeral.

2012 ACM Subject Classification Theory of computation → Type theory; Theory of computation → Proof theory; Theory of computation → Computability

Keywords and phrases cubical type theory, univalence, canonicity, scoping, glueing

Digital Object Identifier 10.4230/LIPIcs.CVIT.2016.23

Funding *Simon Huber*: I acknowledge the support of the Centre for Advanced Study (CAS) in Oslo, Norway, which funded and hosted the research project Homotopy Type Theory and Univalent Foundations during the academic year 2018/19.

Introduction

This paper is a contribution to the analysis of the computational content of the univalence axiom [28] (and higher inductive types). In previous work [2, 3, 4, 5, 20], various *presheaf models* of this axiom have been described in a constructive meta theory. In this formalism, the notion of fibrant type is stated as a refinement of the path lifting operation where one not only provides one of the endpoints but also a partial lift (for a suitable notion of partiality). This generalized form of path lifting operation is a way to state a homotopy extension property, which was recognized very early (see, e.g. [7]) as a key for an abstract development of algebraic topology. The axiom of univalence is then captured by a suitable *equivalence extension operation* (the “glueing” operation), which expresses that we can extend a partially defined equivalence of a given total codomain to a total equivalence. These presheaf models suggest possible extensions of type theory where we manipulate higher dimensional objects [2, 4]. One can define a notion of reduction and prove canonicity for this extension [13]: any closed term of type \mathbb{N} (natural number) is convertible to a numeral. There are however several *non-canonical* choices when defining the path lifting operation by induction on the type, which produce different notion of convertibility.¹ A natural question is how *essential* these non-canonical choices are: can it be that a closed term of type \mathbb{N} , defined without use of such non-canonical reduction rules, becomes convertible to 0 for one choice and 1 for another?

¹ For instance, the definition of this operation for “glue” types is different in [4] and [20].



$\Pi(A, B)'(w)$	$=$	$\Pi(u : A)\Pi(u' : A'u).B'uu'(\mathbf{app}(w, u))$
$\Sigma(A, B)'(w)$	$=$	$\Sigma(u' : A'(w.1)).B'uu'(w.2)$
$\mathbf{fill}_{\psi, b}(u)'$	$=$	$\mathbf{fill}_{\psi, b}(u')$
$\mathbf{Path}(A, a_0, a_1)'(w)$	$=$	$\mathbf{Path}_{\lambda i. A' i(\mathbf{ap}(w, i))} a'_0 a'_1$
$\mathbf{Glue}(A, \psi \mapsto (B, w))'v$	$=$	$\mathbf{Glue}(A'(\mathbf{app}(\mathbf{unglue}, v))) [\psi \mapsto (B'v, (w'.1, \dots))]$

■ **Figure 1** The main rules for the sconing model

The main result of this paper, the *homotopy canonicity theorem*, implies that this cannot be the case. the value of a term is independent of these non-canonical choices. Homotopy canonicity states that, even without providing reduction rules for path lifting operations at type formers, we *still* have that any closed term of type \mathbb{N} is *path equal* to a numeral. (We cannot hope to have convertibility anymore with these path lifting constant.) We can then see this numeral as the “value” of the given term.

Our proof of the homotopy canonicity can be seen as a proof-relevant extension of the *reducibility* or *computability* method, going back to the work of Gödel [11] and Tait [26]. It is however best expressed in an *algebraic* setting. We first define a general notion of model, called *cubical category with families*, defined as a category with families [6] with certain special operations internal to presheaves over a category \mathcal{C} (such as a cube category) with respect to the parameters of an *interval* \mathbb{I} and an object of *cofibrant propositions* \mathbb{F} .

We describe the initial model and how to re-interpret the cubical presheaf models as cubical categories with families. The computability method can then be expressed as a general operation (called “soning”) which applied to an arbitrary model \mathcal{M} produces a new model \mathcal{M}^* with a strict morphism $\mathcal{M}^* \rightarrow \mathcal{M}$. Homotopy canonicity is obtained by applying this general operation to the term model. This construction associates to a (for simplicity, closed) type A a predicate A' on the closed terms $|A|$, and each closed term u a proof u' of $A'u$. The main rules in the closed case are summarized in Figure 1.

Some extensions and variations are then described:

- Our development extends uniformly to identity types and higher inductive types (using the methods of [5]) (Sections 5.1 and 5.2).
- Our development applies equally to the case where one treats univalence instead of glue types as primitive (Appendix C.1). We expect that a similar sconing argument (glueing along a global sections functor to simplicial sets) works to establish homotopy canonicity for the initial split univalent simplicial tribe in the setting of Joyal [14].
- A similar sconing argument adapts to canonicity of cubical type theory with computation of filling at type formers, originally proved by [13] (Appendix C.2).
- Assuming excluded middle, a version of the simplicial set model [15] forms an instance of our development, and distributive lattice cubical type theory interprets in it (Appendix D). Using our technique, one may also reprove canonicity for ordinary Martin Löf type theory with inductive families in a reduction-free way.

Shulman [22] proves homotopy canonicity for homotopy type theory with a truncatedness assumption using the sconing technique. This proof was one starting point for the present work. Some of his constructions may be simplified using our techniques, for example the construction of the natural number type in the sconing.

Setting

We work in a constructive set theory (as presented e.g. in [1]) with a sufficiently long cumulative hierarchy of Grothendieck universes. However, our constructions are not specific

to this setting and can be replayed in other constructive metatheories such as extensional type theory. In Appendix D, we assume classical logic for the discussion of models in simplicial sets.

1 Cubical categories with families

We first recall the notion of categories with families (cwf) [6] equipped with Π - and Σ -types, universes and natural number types. This notion can be interpreted in any presheaf model. In a presheaf model however, we can consider new operations. A *cubical cwf* will be such a cwf in a presheaf model with extra operations which make use of an interval presheaf \mathbb{I} and a type of cofibrant propositions \mathbb{F} as introduced in [5, 20].

1.1 Category with families

Categories with families form an algebraic notion of model of type theory. In order to later model universes à la Russell, we define them in a stratified manner where instead of a single presheaf of types, we specify a filtration of presheaves of “small” types.² The length of the filtration is not essential: we have chosen $1 + \omega$ so that we may specify constructions just at the top level.

A *category with families* (cwf) consists of the following data.

- We have a category of *contexts* \mathbf{Con} and *substitutions* $\mathbf{Hom}(\Delta, \Gamma)$ from Δ to Γ in \mathbf{Con} . The identity substitution on Γ in \mathbf{Con} is written id , and the composition of δ in $\mathbf{Hom}(\Theta, \Delta)$ and σ in $\mathbf{Hom}(\Delta, \Gamma)$ is written $\sigma\delta$.
- We have a presheaf \mathbf{Type} of *types* over the category of contexts. The action of σ in $\mathbf{Hom}(\Delta, \Gamma)$ on a type A over Γ is written $A\sigma$. We have a cumulative sequence of subpresheaves \mathbf{Type}_n of *types of level n* of \mathbf{Type} where n is a natural number.
- We have a presheaf \mathbf{Elem} of *elements* over the category of elements of \mathbf{Type} , i.e. a type $\mathbf{Elem}(\Gamma, A)$ for A in $\mathbf{Type}(\Gamma)$ with $a\sigma$ in $\mathbf{Elem}(\Delta, A\sigma)$ for a in $\mathbf{Elem}(\Gamma, A)$ and σ in $\mathbf{Hom}(\Delta, \Gamma)$ satisfying evident laws.
- We have a terminal context 1 , with the unique element of $\mathbf{Hom}(\Gamma, 1)$ written $()$.
- Given A in $\mathbf{Type}(\Gamma)$, we have a *context extension* $\Gamma.A$. There is a *projection* \mathbf{p} in $\mathbf{Hom}(\Gamma.A, \Gamma)$ and a *generic term* \mathbf{q} in $\mathbf{Elem}(\Gamma.A, A\mathbf{p})$. Given σ in $\mathbf{Hom}(\Delta, \Gamma)$, A in $\mathbf{Type}(\Gamma)$, and a in $\mathbf{Elem}(\Delta, A\sigma)$ we have a *substitution extension* (σ, a) in $\mathbf{Hom}(\Delta, \Gamma.A)$. These operations satisfy $\mathbf{p}(\sigma, a) = \sigma$, $\mathbf{q}(\sigma, a) = a$, and $(\mathbf{p}\sigma, \mathbf{q}\sigma) = \sigma$. Thus, every element of $\mathbf{Hom}(\Delta, \Gamma.A)$ is uniquely of the form (σ, a) with σ and a as above.

We introduce some shorthand notation related to substitution. Given σ in $\mathbf{Hom}(\Delta, \Gamma)$ and A in $\mathbf{Type}(\Gamma)$, we write $\sigma^+ = (\sigma\mathbf{p}, \mathbf{q})$ in $\mathbf{Hom}(\Delta.A\sigma, \Gamma.A)$. Given a in $\mathbf{Elem}(\Gamma, A)$, we write $[a] = (\text{id}, a)$ in $\mathbf{Hom}(\Gamma, \Gamma.A)$. Thus, given B in $\mathbf{Type}_n(\Gamma.A)$ and a in $\mathbf{Elem}(\Gamma, A)$, we have $B[a]$ in $\mathbf{Type}(\Gamma)$. Given furthermore b in $\mathbf{Elem}(\Gamma.A, B)$, we have $b[a]$ in $\mathbf{Elem}(\Gamma, B[a])$. We extend this notation to several arguments: given a_i in $\mathbf{Elem}(\Gamma, A_i)$ for $1 \leq i \leq k$, we write $[a_1, \dots, a_k]$ for $[a_k][a_{k-1}\mathbf{p}] \cdots [a_1\mathbf{p} \dots \mathbf{p}]$ in $\mathbf{Hom}(\Gamma, \Gamma.A_1 \dots A_k)$.

Given a cwf as above, we define what it means to have the following type formers. In addition to the specified laws, all specified operations are furthermore required to be stable under substitution in the evident manner.

² We note that this non-algebraic aspect of the definition does not interfere with the otherwise algebraic character and that subsets and the equality of sets $\mathbf{Elem}(\Gamma, \mathcal{U}_n) = \mathbf{Type}_n$ below could in principle be replaced by injections and natural isomorphisms, respectively.

- **Dependent products.** For A in $\text{Type}(\Gamma)$ and B in $\text{Type}(\Gamma.A)$, we have $\Pi(A, B)$ in $\text{Type}(\Gamma)$, of level n if A and B are. Given b in $\text{Elem}(\Gamma.A, B)$, we have the *abstraction* $\lambda(b)$ in $\text{Elem}(\Gamma, \Pi(A, B))$. Given c in $\text{Elem}(\Gamma, \Pi(A, B))$ and a in $\text{Elem}(\Gamma, A)$, we have the *application* $\text{app}(c, a)$ in $\text{Elem}(\Gamma, B[a])$. These operations satisfy $\text{app}(\lambda(b), a) = b[a]$ and $\lambda(\text{app}(c, q)) = c$.
Given A and B in $\text{Type}(\Gamma)$ we write $A \rightarrow B$ for $\Pi(A, B\mathbf{p})$.
- **Dependent sums.** For A in $\text{Type}(\Gamma)$ and B in $\text{Type}(\Gamma.A)$, we have $\Sigma(A, B)$ in $\text{Type}(\Gamma)$, of level n if A and B are. Given a in $\text{Elem}(\Gamma, A)$ and b in $\text{Elem}(\Gamma, B[a])$, we have the *pairing* $\text{pair}(a, b)$ in $\text{Elem}(\Gamma, \Sigma(A, B))$. Given c in $\text{Elem}(\Gamma, \Sigma(A, B))$, we have the *first projection* $\text{fst}(c)$ in $\text{Elem}(\Gamma, A)$ and *second projection* $\text{snd}(c)$ in $\text{Elem}(\Gamma, B[\text{fst}(c)])$. These operations satisfy $\text{fst}(\text{pair}(a, b)) = a$, $\text{snd}(\text{pair}(a, b)) = b$, and $\text{pair}(\text{fst}(c), \text{snd}(c)) = c$. Thus, every element of $\text{Elem}(\Gamma, \Sigma(A, B))$ is uniquely of the form $\text{pair}(a, b)$ with a and b as above.
Given A and B in $\text{Type}(\Gamma)$ we write $A \times B$ for $\Sigma(A, B\mathbf{p})$.
- **Universes.** We have \mathbf{U}_n in $\text{Type}_{n+1}(\Gamma)$ such that $\text{Type}_n(\Gamma) = \text{Elem}(\Gamma, \mathbf{U}_n)$, and the action of substitutions on $\text{Elem}(\Gamma, \mathbf{U}_n)$ is compatible with that on $\text{Type}_n(\Gamma)$.
- **Natural numbers.** We have \mathbf{N} in $\text{Type}(\Gamma)$ with *zero* 0 in $\text{Elem}(\Gamma, \mathbf{N})$ and *successor* $\mathbf{S}(n)$ in $\text{Elem}(\Gamma, \mathbf{N})$ for n in $\text{Elem}(\Gamma, \mathbf{N})$. Given P in $\text{Type}(\Gamma.\mathbf{N})$, z in $\text{Elem}(\Gamma, P[0])$, s in $\text{Elem}(\Gamma.\mathbf{N}.P, P(\mathbf{p}, \mathbf{S}(\mathbf{q}))\mathbf{p})$, and $n : \text{Elem}(\Gamma, \mathbf{N})$, we have the *elimination* $\text{natrec}(P, z, s, n)$ in $\text{Elem}(\Gamma, P[n])$, satisfying as usual $\text{natrec}(P, z, s, 0) = z$ and $\text{natrec}(P, z, s, \mathbf{S}(n)) = s[n, \text{natrec}(P, z, s, n)]$.

A *structured cwf* is a cwf with type formers as above.

A (strict) morphism $\mathcal{M} \rightarrow \mathcal{N}$ of cwfs is defined in the evident manner and consists of a functor $F : \text{Con}_{\mathcal{M}} \rightarrow \text{Con}_{\mathcal{N}}$ and natural transformations $u : \text{Type}_{\mathcal{M}} \rightarrow \text{Type}_{\mathcal{N}}F$ and $v : \text{Elem}_{\mathcal{M}} \rightarrow \text{Elem}_{\mathcal{N}}(F, u)$ such that v restricts to types of level n and the terminal context and context extension is preserved strictly. A morphism $\mathcal{M} \rightarrow \mathcal{N}$ of structured cwfs additionally preserves the operations of the above type formers. We obtain a category of structured cwfs.

1.2 Internal language of presheaves

For the rest of the paper, we fix a category \mathcal{C} in the lowest Grothendieck universe. As in [2, 20, 18], we will use the language of extensional type theory (with subtypes) to describe constructions in the presheaf topos over \mathcal{C} .

In the interpretation of this language, a context is a presheaf A over \mathcal{C} , a type B over A is a presheaf over the category of elements of A , and an element of B is a section. A *global type* is a type in the global context, i.e. a presheaf over \mathcal{C} . Similarly, a *global element* of a global type is a section of that presheaf.

Given a dependent type B over a type A , we think of B as a family of types $B a$ indexed by elements a of A . We have the usual dependent sum $\Sigma(a : A).B(a)$ and dependent product $\Pi(a : A).B a$, with projections of $s : \Sigma(a : A).B a$ written $s.1$ and $s.2$ and application of $f : \Pi(a : A).B a$ to $a : A$ written $f a$. We also use the categorical pairing $\langle f, g \rangle : X \rightarrow \Sigma(a : A).B$ given $f : X \rightarrow A$ and $g : \Pi(x : X).B(f x)$ and other commonly used notations. The hierarchy of Grothendieck universes in the ambient set theory gives rise to a cumulative hierarchy $\mathbf{U}_0, \mathbf{U}_1, \dots, \mathbf{U}_\omega$ of universes à la Russell. We model propositions as subtypes of a fixed type $\mathbf{1}$ with unique element tt . We have subuniverses $\Omega_i \subseteq \mathbf{U}_i$ of propositions for $i \in \{0, 1, \dots, \omega\}$.

When working in this internal language, we refer to the types as “sets” to avoid ambiguity with the types of (internal) cwfs we will be considering.

1.3 Cubical categories with families

We now work internally to presheaves over \mathcal{C} . We assume the following:

- an *interval* $\mathbb{I} : \mathbb{U}_0$ with *endpoints* $0, 1 : \mathbb{I}$,
- an object $\mathbb{F} : \mathbb{U}_0$ of *cofibrant propositions* with a monomorphism $[-] : \mathbb{F} \rightarrow \Omega_0$.

As in [5, 20], a *partial element* of a set T is given by an element φ in \mathbb{F} and a function $[\varphi] \rightarrow T$. We say that a total element v of T extends such a partial element φ, u if we have $[\varphi] \rightarrow u \text{tt} = v$.

Given $A : \mathbb{I} \rightarrow \mathbb{U}_\omega$, we write $\text{hasFill}(A)$ for the set of operations taking as inputs φ in \mathbb{F} , $b \in \{0, 1\}$, and a partial section u in $\Pi(i : \mathbb{I}).[\varphi] \vee (i = b) \rightarrow A i$ and producing an extension of u to a total section in $\Pi(i : \mathbb{I}) A i$. Given a set X and $Y : X \rightarrow \mathbb{U}_\omega$, we write $\text{Fill}(X, Y)$ for the set of *filling structures* on Y , producing an element of $\text{hasFill}(Y \circ x)$ for x in $\mathbb{I} \rightarrow X$. Given s in $\text{Fill}(X, Y)$ and x, φ, b, u as above, we write $s(x, \varphi, b u)$ for the resulting total section in $\Pi(i : \mathbb{I}).Y(x i)$.

We now interpret the definitions of Section 1.1 in the internal language of the presheaf topos. A *cubical cwf* is a structured cwf denoted as before that additionally has the following *cubical operations and type formers*. Again, all specified operations are required to be stable under substitution.

- **Filling operation.** We have fill in $\text{Fill}(\text{Type}(\Gamma), \lambda A. \text{Elem}(\Gamma, A))$ for Γ in Con . Let us spell out stability under substitution: given $A : \mathbb{I} \rightarrow \text{Type}(\Gamma)$, φ in \mathbb{F} , $b \in \{0, 1\}$, u in $\Pi(i : \mathbb{I}).[\varphi] \vee (i = b) \rightarrow \text{Elem}(\Gamma, A i)$, and σ in $\text{Hom}(\Delta, \Gamma)$ and $r : \mathbb{I}$, we have $(\text{fill}(A, \varphi, b, u) r) \sigma = \text{fill}(\lambda i. (A i) \sigma, \varphi, b, \lambda i x. (u i x) \sigma) r$.

Note that we do not include computation rules for fill at type formers. This corresponds to our decision to treat fill as a non-canonical operation.

- **Dependent path types.** Given A in $\mathbb{I} \rightarrow \text{Type}(\Gamma)$ with a_b in $\text{Elem}(\Gamma, A b)$ for $b \in \{0, 1\}$, we have a type $\text{Path}(A, a_0, a_1)$ in $\text{Type}(\Gamma)$, of level n if A is. Given u in $\Pi(i : \mathbb{I}).\text{Elem}(\Gamma, A i)$, we have the *path abstraction* $\langle \rangle(u)$ in $\text{Elem}(\Gamma, \text{Path}(A, u 0, u 1))$. Given p in $\text{Elem}(\Gamma, \text{Path}(A, a_0, a_1))$ and i in \mathbb{I} , we have the *path application* $\text{ap}(p, r)$ in $\text{Elem}(\Gamma, A i)$. These operations satisfy the laws $\text{ap}(p, b) = a_b$ for $b \in \{0, 1\}$, $\text{ap}(\langle \rangle(u), i) = u i$, and $\langle \rangle(\lambda i. \text{ap}(p, i)) = p$. Thus, every element of $\text{Elem}(\Gamma, \text{Path}(A, a_0, a_1))$ is uniquely of the form $\langle \rangle(u)$ with u in $\Pi(i : \mathbb{I}).\text{Elem}(\Gamma, A i)$ such that $u 0 = a_0$ and $u 1 = a_1$.

Using path types, we define $\text{isContr}(A)$ in $\text{Type}(\Gamma)$ for A in $\text{Type}(\Gamma)$ as well as isEquiv in $\text{Type}(\Gamma. A \rightarrow B)$ and Equiv in $\text{Type}(\Gamma)$ for A, B in $\text{Type}(\Gamma)$ as in [4]. These notions are used in the following type former, which extends any partially defined equivalence (given total codomain) to a totally defined function.

- **Glue types.** Given A in $\text{Type}(\Gamma)$, φ in \mathbb{F} , T in $[\varphi] \rightarrow \text{Type}(\Gamma)$ and $e : [\varphi] \rightarrow \text{Elem}(\Gamma, \text{Equiv}(T \text{tt}, A))$, we have the *glueing* $\text{Glue}(A, \varphi, T, e)$ in $\text{Type}(\Gamma)$, equal to T on $[\varphi]$ and of level n if A and T are. We have unglue in $\text{Elem}(\Gamma, \text{Glue}(A, \varphi, T, e) \rightarrow A)$ such that $\text{unglue} = \text{fst}(e) \text{tt}$ on $[\varphi]$. Given a in $\text{Elem}(\Gamma, A)$ and t in $[\varphi] \rightarrow \text{Elem}(\Gamma, T)$ such that $\text{app}(\text{fst}(e) \text{tt}, t \text{tt}) = a$ on $[\varphi]$, we have $\text{glue}(a, t)$ in $\text{Elem}(\Gamma, \text{Glue}(A, \varphi, T, e))$. These operations satisfy $\text{app}(\text{unglue}, \text{glue}(a, t)) = a$ and $\text{glue}(\text{app}(\text{unglue}, u), \lambda x. u) = u$. Thus, every element of $\text{Elem}(\Gamma, \text{Glue}(A, \varphi, T, e))$ is uniquely of the form $\text{glue}(a, t)$ with a and t as above.

The notion of morphism of structured cwfs lifts to an evident notion of morphism of cubical cwfs. We obtain, internally to presheaves over \mathcal{C} , a category of cubical cwfs. We now lift this category of cubical cwfs from the internal language to the ambient theory by interpreting it in the global context: externally, a cubical cwf (relative to the chosen base

category \mathcal{C} , interval \mathbb{I} , and cofibrant propositions \mathbb{F}) consists of a presheaf Con over \mathcal{C} , a presheaf Type over the category of elements of Con , etc.

► **Remark 1.** Fix a cubical cwf as above. Assume that \mathbb{I} has a connection algebra structure and that \mathbb{F} forms a sublattice of Ω_0 that contains the interval endpoint inclusions. As in [4], it is then possible in the above context of the glue type former to construct an element of $\text{Elem}(\Gamma, \text{iEquip}[\text{unglue}])$. From this, one derives an element of $\text{Elem}(\Gamma, \text{iUnivalence}_n)$ where $\text{iUnivalence}_n = \Pi(\mathbb{U}_n, \text{iContr}(\Sigma(\mathbb{U}_n, \text{Equiv}(\mathbf{q}, \mathbf{qp}))))$ for $n \geq 0$, i.e. univalence is provable. One may also show that the path type applied to constant families $\mathbb{I} \rightarrow \text{Type}(\Gamma)$ interprets the rules of identity types of Martin-Löf with the computation rule for the eliminator J replaced by a propositional equality. Thus, we obtain an interpretation of univalent type theory with identity types with propositional computation in any cubical cwf.

2 Two examples of cubical cwfs

In this section we give two examples of cubical cwfs: a term model and a particular cubical cwf formulated in a constructive metatheory, the latter with extra assumptions on \mathbb{I} and \mathbb{F} .

2.1 Term model

We sketch how to give a cubical cwf \mathcal{T} built from syntax, and refer the reader to Appendix A for more details. All our judgments will be indexed by an object X of \mathcal{C} and given a judgment $\Gamma \vdash_X \mathcal{J}$ and $f: Y \rightarrow X$ in \mathcal{C} we get $\Gamma f \vdash_Y \mathcal{J}f$. Here, f acts on expressions as an implicit substitution, while for substitutions on object variables we will use explicit substitutions.

The forms of judgment are:

$$\Gamma \vdash_X \quad \Gamma \vdash_X A \quad \Gamma \vdash_X A = B \quad \Gamma \vdash_X t : A \quad \Gamma \vdash_X t = u : A \quad \sigma : \Delta \rightarrow_X \Gamma$$

The main rules are given in the appendix. This then induces a cubical cwf \mathcal{T} by taking, say, the presheaf of contexts at stage X to be equivalence classes of Γ for $\Gamma \vdash_X$ where the equivalence relation is judgmental equality.

Some rules are a priori infinitary, but in some cases (such as the one considered in [4]) it is possible to present the rules in a finitary way.

This formal system expresses the laws of cubical cwfs in rule form, and thus can be interpreted in an arbitrary cubical cwf in the usual way:

► **Theorem 2.** *Fixing parameters $\mathcal{C}, \mathbb{I}, \mathbb{F}$, we have the cubical cwf \mathcal{T} initial cubical in the category of cubical cwfs.*

2.2 Developments in presheaves over \mathcal{C}

We now assume that \mathbb{I} and \mathbb{F} satisfy the axioms $\text{ax}_1, \dots, \text{ax}_9$ of [20], internally to presheaves over \mathcal{C} . We also make an external assumption, namely that the endofunctor on presheaves over \mathcal{C} of exponentiation with \mathbb{I} has a right adjoint R that preserves global types of level n . This is e.g. the case if \mathbb{I} is representable and \mathcal{C} is closed under finite products.

Most of the arguments will be done in the internal language of the presheaf topos. At certain points however, we need to consider the set of global sections of a global type F ; we denote this by $\square F$. We stress that statements involving \square are external, not to be interpreted in the internal language. Crucially, the adjunction $(-)^{\mathbb{I}} \dashv R$ cannot be made internal [18].

Recall the global type $\text{hasFill} : \mathbb{U}_\omega^{\mathbb{I}} \rightarrow \mathbb{U}_\omega$ from Section 1.3. Taking the slice over \mathbb{U}_ω , the adjunction $(-)^{\mathbb{I}} \dashv R$ descends to an adjunction between categories of types over \mathbb{U}_ω and

$\mathbb{U}_\omega^{\mathbb{I}}$. Applying the right adjoint of this adjunction to hasFill , we obtain global $\mathbf{C} : \mathbb{U}_\omega \rightarrow \mathbb{U}_\omega$ such that naturally in a global type X with global $Y : X \rightarrow \mathbb{U}_\omega$, global elements of $(x : X^{\mathbb{I}}) \rightarrow \text{hasFill}(Y \circ x)$ are in bijection with global elements of $(x : X) \rightarrow \mathbf{C}(Y x)$. Given a global type X and global $Y : X \rightarrow \mathbb{U}_\omega$, we thus have a logical equivalence (maps back and forth)

$$\square \text{Fill}(X, Y) \longleftrightarrow \square \Pi(x : X) \mathbf{C}(Y x) \quad (1)$$

natural in X .³ Note that \mathbf{C} descends to $\mathbf{C} : \mathbb{U}_n \rightarrow \mathbb{U}_n$ for $n \geq 0$. We write $\mathbb{U}_i^{\text{fib}} = \Sigma(A : \mathbb{U}_i) \mathbf{C}(A)$ for $i \in \{0, 1, \dots, \omega\}$; we call $\mathbb{U}_i^{\text{fib}}$ a universe of *fibrant sets*. Now set $X = \mathbb{U}_\omega^{\text{fib}}$ and $Y(A, c) = A$ in (1). We trivially have $\square \Pi(x : X) \mathbf{C}(Y x)$, thus get

$$\text{fill} : \text{Fill}(\mathbb{U}_\omega^{\text{fib}}, \lambda(A, c).A). \quad (2)$$

This is essentially the counit of the adjunction defining \mathbf{C} . Note that [18] use modal extensions of type theory to perform this reasoning internal to presheaves over \mathcal{C} .

► **Remark 3.** Internally, a map $\text{Fill}(X, Y) \rightarrow \Pi(x : X) \mathbf{C}(Y x)$ does not generally exist for a set X and $Y : X \rightarrow \mathbb{U}_\omega$ as for $X = 1$ one would derive a filling structure for any “homogeneously fibrant” set, which is impossible (see [20, Remark 5.9]). However, from (2) we get a map $\Pi(x : X) \mathbf{C}(Y x) \rightarrow \text{Fill}(X, Y)$ natural in X using closure of filling structures under substitution (see below).

We recall some constructions of [4, 20] in the internal language.

- Given $A : \mathbb{I} \rightarrow \mathbb{U}_\omega$ and $a_b : A_b$ for $b \in \{0, 1\}$, *dependent paths* $\text{Path}_A a_0 a_1$ are the set of maps $p : \Pi(i : \mathbb{I}).A i$ such that $p 0 = a_0$ and $p 1 = a_1$. We use the same notation for non-dependent paths.
- For $A : \mathbb{U}_\omega$, we have a set $\text{isContr}(A)$ of witnesses of *contractibility*, defined using paths.
- Given $A, B : \mathbb{U}_\omega$ with $f : A \rightarrow B$, we have the set $\text{isEquiv}(f)$ with elements witnessing that f is an *equivalence*, defined using contractibility of homotopy fibers. We write $\text{Equiv}(A, B) = \Sigma(f : A \rightarrow B).\text{isEquiv}(f)$.
- Given $A : \mathbb{U}_\omega$, $\varphi : \mathbb{F}$, $B : [\varphi] \rightarrow \mathbb{U}_\omega$, and $e : [\varphi] \rightarrow \text{Equiv}(B \text{tt}, A)$, the *glueing* $\text{Glue } A [\varphi \mapsto (B, e)]$ consists of elements $\text{glue } a [\varphi \mapsto b]$ with $a : A$ and $b : [\varphi] \rightarrow B$ such that $e.1 (b \text{tt}) = a$ on $[\varphi]$ and is defined in such a way that $\text{Glue } A [\varphi \mapsto (B, e)] = T \text{tt}$ and $\text{glue } a [\varphi \mapsto b] = b \text{tt}$ on $[\varphi]$. The canonical map $\text{unglue} : \text{Glue } A [\varphi \mapsto (B, e)] \rightarrow A$ is an equivalence.

These operations are valued in \mathbb{U}_n if their inputs are. We further recall basic facts from [20] about filling structures in the internal language.

- Filling structures are closed under substitution: given $f : X' \rightarrow X$ and $Y : X \rightarrow \mathbb{U}_\omega$, any element of $\text{Fill}(X, Y)$ induces an element of $\text{Fill}(X', Y \circ f)$, naturally in X' .
- Filling structures are closed under exponentiation: given sets S, X and $Y : X \rightarrow \mathbb{U}_\omega$, any element of $\text{Fill}(X, Y)$ induces an element of $\text{Fill}(X^S, \lambda x.\Pi(s : S).Y(x s))$, naturally in S .
- Filling structures are closed under $\Pi, \Sigma, \text{Path}, \text{Glue}$. E.g. for dependent products, given $A : \Gamma \rightarrow \mathbb{U}_\omega$ with $\text{Fill}(\Gamma, A)$ and $B : \Pi(\rho : \Gamma).A \rho \rightarrow \mathbb{U}_\omega$ with $\text{Fill}(\Sigma(\rho : \Gamma).A \rho, \lambda(\rho, a).B \rho a)$, we have $\text{Fill}(\Gamma, \lambda(\rho : \Gamma).\Pi(a : A \rho).B \rho a)$.

From the last point, we deduce using that \mathbf{C} is closed under $\Pi, \Sigma, \text{Path}, \text{Glue}$, and that $\mathbf{C}(A)$ implies $\mathbf{C}(A^S)$ for $A, S : \mathbb{U}_\omega$.⁴ Let us explain this in the case of dependent products. We apply (1) with a suitable “generic context” $X = \Sigma((A, c) : \mathbb{U}_\omega^{\text{fib}}).A \rightarrow \mathbb{U}_\omega^{\text{fib}}$ and

³ We record only the logical equivalence instead of an isomorphism so that it will be easier to apply our constructions in situations where the right adjoint R fails to exist, see Appendix D. Naturality is only used at a one point below, for the forward map, to construct suitable elements of \mathbf{C} applied to glueings.

⁴ Note that naturality in S of the latter operation is used in substitutional stability of universes in the scoping in Section 3.

$Y((A, c), \langle B, d \rangle) = \Pi(a : A).B a$. Using the map of Remark 3 and closure of filling structures under dependent product, we have $\square \text{Fill}(X, Y)$ and can conclude $\mathbf{C}(\Pi(a : A).B a)$ for $(A, c) : \mathbf{U}_\omega^{\text{fib}}$ and $\langle B, d \rangle : A \rightarrow \mathbf{U}_\omega^{\text{fib}}$. Note that in the case of **Glue** with $(A, c) : \mathbf{U}_\omega^{\text{fib}}$, $\varphi : \mathbb{F}$, $\langle B, d \rangle : [\varphi] \rightarrow \mathbf{U}_\omega^{\text{fib}}$, and $e : [\varphi] \rightarrow \mathbf{Equiv}(B \text{tt}, A)$, naturality of the forward map of (1) is needed to see that the element $c : \mathbf{C}(\text{Glue } A [\varphi \mapsto (B, e)])$ constructed in the same fashion as above for dependent products equals $d \text{tt} : \mathbf{C}(B \text{tt})$ on $[\varphi]$.

As in [4, 20, 18], glueing shows $\text{Fill}(1, \mathbf{U}_n^{\text{fib}})$ for $n \geq 0$. Using (1), we conclude $\mathbf{C}(\mathbf{U}_n^{\text{fib}})$.

Let \mathbb{N} denote the natural number object in presheaves over \mathcal{C} . From [4, 20], we have $\text{Fill}(1, \mathbb{N})$. Using (1), we conclude $\mathbf{C}(\mathbb{N})$.

We justify fibrant indexed inductive sets in Appendix B.

2.3 Standard model

Making the same assumptions on $\mathcal{C}, \mathbb{I}, \mathbb{F}$ as in Section 2.2, we can now specify the standard model \mathcal{S} of cubical type theory in the sense of the current paper as a cubical cwf (with respect to parameters $\mathcal{C}, \mathbb{I}, \mathbb{F}$) purely using the internal language of the presheaf topos. The cwf is induced by the family over $\mathbf{U}_\omega^{\text{fib}}$ given by the first projection as follows.

- The category of contexts is \mathbf{U}_ω , with $\text{Hom}(\Delta, \Gamma)$ the functions from Δ to Γ .
- The types over Γ are maps from Γ to $\mathbf{U}_\omega^{\text{fib}}$; a type $\langle A, p \rangle$ is of level n if A is in $\Gamma \rightarrow \mathbf{U}_n$. This is clearly functorial in Γ .
- The elements of $\langle A, p \rangle : \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$ are $\Pi(\rho : \Gamma).A \rho$. This is clearly functorial in Γ .
- The terminal context is given by 1.
- The context extension of Γ by $\langle A, p \rangle$ is given by $\Sigma(\rho : \Gamma).A \rho$, with \mathbf{p}, \mathbf{q} given by projections and substitution extension given by pairing.

We briefly go through the necessary type formers and operations, omitting evident details.

- The dependent product of $\langle A, c \rangle : \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$ and $\langle B, d \rangle : \Sigma(\rho : \Gamma).A \rho \rightarrow \mathbf{U}_\omega^{\text{fib}}$ is $\langle \lambda \rho. \Pi(a : A \rho).B(\rho, a), e \rangle$ where $e \rho : \mathbf{C}(\Pi(a : A \rho).B(\rho, a))$ is induced by $c \rho : \mathbf{C}(A \rho)$ and $d \rho a : \mathbf{C}(B(\rho, a))$ for $a : A$ as discussed above.
- The dependent sum of $\langle A, c \rangle : \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$ and $\langle B, d \rangle : \Sigma(\rho : \Gamma).A \rho \rightarrow \mathbf{U}_\omega^{\text{fib}}$ is $\langle \lambda \rho. \Sigma(a : A \rho).B(\rho, a), e \rangle$ where e is induced by c and d .
- The universe $\mathbf{U}_n : \Gamma \rightarrow \mathbf{U}_{n+1}^{\text{fib}}$ is constantly $(\mathbf{U}_n^{\text{fib}}, c)$ with $c : \mathbf{C}(\mathbf{U}_n^{\text{fib}})$ as above.
- The natural number type $\mathbb{N} : \Gamma \rightarrow \mathbf{U}_0^{\text{fib}}$ is constantly (\mathbb{N}, c) with $c : \mathbf{C}(\mathbb{N})$ as above. The zero and successor constructors and eliminator are given by the corresponding features of the natural number object \mathbb{N} .

We now turn to the cubical aspects.

- The filling operation $\text{fill} : \text{Fill}(\Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}, \lambda \langle A, p \rangle. \Pi(\rho : \Gamma).A \rho)$ is derived from (2) by closure of filling structures under exponentiation.
- Given $\langle A, c \rangle : \mathbb{I} \rightarrow \Gamma \rightarrow \Sigma(A : \mathbf{U}_\omega) \mathbf{C}(A)$ and $a_b : \Pi(\rho : \Gamma).A b \rho$ for $b \in \{0, 1\}$, we define $\text{Path}(A, a_0, a_1) : \Gamma \rightarrow \Sigma(A : \mathbf{U}_\omega) \mathbf{C}(A)$ as $\langle \Pi(\rho : \Gamma). \text{Path}_{\lambda i. A i \rho} c_0 c_1 \rangle, d \rangle$ where $d \rho : \mathbf{C}(\text{Path}_{\lambda i. A i \rho} c_0 c_1)$ induced by $\lambda i. c i \rho$.

Before defining glue types, we note that the notions **isContr** and **isEquiv** in the cubical cwf we are defining correspond to the notions **isContr** and **isEquiv**. For example, given a type $A : \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$, then the elements of **isContr**(A), given by $\Pi(\rho : \Gamma). \text{isContr}(A).1 \rho$, are in bijection with $\Pi(\rho : \Gamma). \text{isContr}(A).1 \rho$ naturally in Γ .

- Given $\langle A, c \rangle : \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$, $\varphi : \mathbb{F}$, $\langle T, d \rangle : [\varphi] \rightarrow \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$ and $e : [\varphi] \rightarrow \mathbf{Equiv}(T \text{tt}, A)$, we define $\text{Glue}(\langle A, c \rangle, \varphi, \langle T, d \rangle, e) : \Gamma \rightarrow \mathbf{U}_\omega^{\text{fib}}$ as $\lambda \rho. (\text{Glue}(A \rho)[\varphi \mapsto (T \text{tt } \rho, e' \text{tt } \rho)], q \rho)$ where $e' \text{tt } \rho : \mathbf{Equiv}(T \text{tt } \rho, A \rho)$ is induced by $e \text{tt } \rho$ and $q \rho$ is induced by $c \rho$ and $\lambda x. d x \rho$.

We have thus verified the following statement.

► **Theorem 4.** *Assuming the parameters $\mathcal{C}, \mathbb{I}, \mathbb{F}$ satisfy the assumptions of Section 2.2, the standard model \mathcal{S} forms a cubical cwf.*

3 Scoring

We make the same assumptions on our parameters $\mathcal{C}, \mathbb{I}, \mathbb{F}$ as in Section 2.2. Given a cubical cwf \mathcal{M} (with respect to these parameters) denoted $\text{Con}, \text{Hom}, \dots$ as in Section 1.3, we will define a new cubical cwf \mathcal{M}^* denoted $\text{Con}^*, \text{Hom}^*, \dots$, the (Artin) glueing of \mathcal{M} with the standard model \mathcal{S} along an (internal) global sections functor, i.e. the *scoring* of \mathcal{M} .

Recall from Section 1.3 the operation fill of \mathcal{M} . Instantiating it to the terminal context, we get $\square\text{Fill}(\text{Type}(1), \lambda A. \text{Elem}(1, A))$. Using the forward direction of Equation (1), we thus have an internal operation $k : \Pi(A : \text{Type}(1)) \rightarrow \mathbb{C}(\text{Elem}(1, A))$.

From now on, we will work in the internal language of presheaves over \mathcal{C} . We start by defining a global sections operation $|-|$ mapping contexts, types, and elements of \mathcal{M} to those of \mathcal{S} .

- Given $\Gamma : \text{Con}$, we define $|\Gamma| : \mathbb{U}_\omega$ as the set of substitutions $\text{Hom}(1, \Gamma)$. Given a substitution $\sigma : \text{Hom}(\Delta, \Gamma)$, we define $|\sigma| : |\Delta| \rightarrow |\Gamma|$ as $|\sigma|\rho = \sigma\rho$. This evidently defines a functor.
- Given $A : \text{Type}(\Gamma)$, we define $|A| : |\Gamma| \rightarrow \mathbb{U}_\omega^{\text{fib}}$ as $|A|\rho = (\text{Elem}(1, A\rho), k(A\rho))$. This evidently natural in Γ . If A is of level n , then $|A| : |\Gamma| \rightarrow \mathbb{U}_n^{\text{fib}}$.
- Given $a : \text{Elem}(\Gamma, A)$ we define $|a| : \Pi(\rho : \Gamma). (|A|\rho).1$ as $|a|\rho = a\rho$. This is evidently natural in Γ .

Note that $|-|$ preserves the terminal context and context extension up to canonical isomorphism in the category of contexts. One could thus call $|-|$ an (internal) *pseudomorphism* cwf's from \mathcal{M} to \mathcal{S} . The scoring \mathcal{M}^* will be defined as essentially the (Artin) glueing along this pseudomorphism, but we will be as explicit as possible and not define (Artin) glueing at the level of generality of an abstract pseudomorphism.

For convenience, we also just write $|A| : |\Gamma| \rightarrow \mathbb{U}_\omega$ instead of $\lambda\rho. (|A|\rho).1$, implicitly applying the first projection. We also write just $|A|$ for $|A|(|\cdot|)$ if Γ is the terminal context.

3.1 Contexts, substitutions, types and elements

We start by defining the cwf \mathcal{M}^* .

- A context $(\Gamma, \Gamma') : \text{Con}^*$ consists of a context $\Gamma : \text{Con}$ in \mathcal{M} and a family Γ' over $|\Gamma|$ (which in the context of Artin glueing should be thought of as a substitution in \mathcal{S} from some context to $|\Gamma|$). We think of Γ' as a *proof-relevant computability predicate*. A substitution $(\sigma, \sigma') : \text{Hom}^*((\Delta, \Delta'), (\Gamma, \Gamma'))$ consists of a substitution $\sigma : \Delta \rightarrow \Gamma$ in \mathcal{M} and a map $\sigma' : \Pi(\nu : |\Delta|). \Delta'(\nu) \rightarrow \Gamma'(\sigma\nu)$. This evidently has the structure of a category.
- A type $(A, A') : \text{Type}^*(\Gamma, \Gamma')$ consists of a type $A : \text{Type}(\Gamma)$ in \mathcal{M} and

$$A' : \Pi(\rho : |\Gamma|)(\rho' : \Gamma'\rho). |A|\rho \rightarrow \mathbb{U}_\omega^{\text{fib}}.$$

We think of A' as a fibrant *proof-relevant computability family* on A . In the abstract context of Artin glueing for cwf's, we should think of it as an element of $\text{Type}(\Sigma(\rho : |\Gamma|)(\rho' : \Gamma'\rho). |A|\rho)$ in \mathcal{S} , but this point of view is not compatible with the universes à la Russell we are going to model. Recalling $\mathbb{U}_\omega^{\text{fib}} = \Sigma(X : \mathbb{U}_\omega). \mathbb{C}(X)$, we also write $\langle A', \text{fib}_{A'} \rangle$ instead of A' if we want to directly access the family and split off its proof of fibrancy.

The type (A, A') is of level n if A and A' are.

23:10 Homotopy canonicity for cubical type theory

The action of a substitution $(\sigma, \sigma') : \text{Hom}^*((\Delta, \Delta'), (\Gamma, \Gamma'))$ on (A, A') is given by

$$(A\sigma, \lambda\nu\nu' a.A'(\sigma\nu)(\sigma'\nu\nu') a).$$

- An element $(a, a') : \text{Elem}^*((\Gamma, \Gamma'), (A, \langle A', \text{fib}_{A'} \rangle))$ consists of $a : \text{Elem}(\Gamma, A)$ in \mathcal{M} and

$$a' : \Pi(\rho : |\Gamma|)(\rho' : \Gamma' \rho). A'(\rho, \rho', a\rho).$$

In the context of Artin glueing (with types in \mathcal{M}^* presented correspondingly), this should be thought of as an element $a' : \text{Elem}(\Sigma(\rho : |\Gamma|). \Gamma' \rho, \lambda(\rho, \rho'). A'(\rho, \rho', |a|\rho))$ of \mathcal{S} .

The action of a substitution $(\sigma, \sigma') : \text{Hom}^*((\Delta, \Delta'), (\Gamma, \Gamma'))$ on the element (a, a') is given by $(a\sigma, \lambda\nu\nu'. a' \sigma\nu(\sigma'\nu\nu'))$.

- The terminal context is given by $(1, 1')$ defined by $1'() = 1$.
- The extension in \mathcal{M}^* of a context (Γ, Γ') by a type (A, A') is given by $(\Gamma.A, (\Gamma.A)')$ where $(\Gamma.A')(\rho, a) = \Sigma(\rho' : \Gamma' \rho). (A' \rho \rho' a). 1$. The projection $\mathbf{p}^* : \text{Hom}^*((\Gamma, \Gamma').(A, A'), (\Gamma, \Gamma'))$ is $(\mathbf{p}, \mathbf{p}')$ where $\mathbf{p}'(\rho, a)(\rho', a') = a'$ and the generic term $\mathbf{q}^* : \text{Elem}((\Gamma, \Gamma').(A, A')\mathbf{p}^*)$ is $(\mathbf{q}, \mathbf{q}')$ where $\mathbf{q}'(\rho, a)(\rho', a') = a'$. The extension of $(\sigma, \sigma') : \text{Hom}^*((\Delta, \Delta'), (\Gamma, \Gamma'))$ with $(a, a') : \text{Elem}^*((\Delta, \Delta'), (A, A')(\sigma, \sigma'))$ is $((\sigma, a), \lambda\nu\nu'. (\sigma'\nu\nu', a'\nu\nu'))$.

3.2 Type formers and operations

3.2.1 Dependent products

Let $(A, \langle A', \text{fib}_{A'} \rangle) : \text{Type}^*(\Gamma, \Gamma')$ and $(B, \langle B', \text{fib}_{B'} \rangle) : (\Gamma, \Gamma').(A, \langle A', \text{fib}_{A'} \rangle)$. We define the dependent product $\Pi^*((A, \langle A', \text{fib}_{A'} \rangle), (B, \langle B', \text{fib}_{B'} \rangle)) = (\Pi(A, B), (\Pi(A, B)', \text{fib}_{\Pi(A, B)'})$ where

$$\Pi(A, B)'(\rho, \rho', f) = \Pi(a : |A|\rho)(a' : A' \rho \rho' a). B'(\rho, a)(\rho', a')(\mathbf{app}(f, a))$$

and $\text{fib}_{\Pi(A, B)' }(\rho, \rho', f)$ is given by closure of \mathbf{C} under dependent product applied to $(|A|\rho). 2$, $\text{fib}_{A'} \rho \rho' a$ for $a : |A|\rho$, and $\text{fib}_{B'}(\rho, a)(\rho', a')(\mathbf{app}(f, a))$ for additionally $a' : A' \rho \rho' a$.

Given an element (b, b') of $(B, \langle B', d \rangle)$ in \mathcal{M}^* , we define the abstraction $\mathbf{lam}^*(b, b') = (\mathbf{lam}(b), \mathbf{lam}(b'))$ where $\mathbf{lam}(b)' \rho \rho' a a' = b'(\rho, a)(\rho', a')$.

Given elements (f, f') of $\Pi^*((A, \langle A', c \rangle), (B, \langle B', d \rangle))$ and (a, a') of $(A, \langle A', \text{fib}_{A'} \rangle)$ in \mathcal{M}^* , we define the application $\mathbf{app}^*((f, f'), (a, a')) = (\mathbf{app}(f, a), \mathbf{app}(f, a'))$ where $\mathbf{app}(f, a)' \rho \rho' = f' \rho \rho' a \rho (a' \rho \rho')$.

3.2.2 Dependent sums

Let $(A, \langle A', \text{fib}_{A'} \rangle) : \text{Type}^*(\Gamma, \Gamma')$ and $(B, \langle B', \text{fib}_{B'} \rangle) : (\Gamma, \Gamma').(A, \langle A', \text{fib}_{A'} \rangle)$. We define the dependent sum $\Sigma^*((A, \langle A', \text{fib}_{A'} \rangle), (B, \langle B', \text{fib}_{B'} \rangle)) = (\Sigma(A, B), \langle \Sigma(A, B)', \text{fib}_{\Sigma(A, B)' } \rangle)$ where

$$\Sigma(A, B)' \rho \rho' (\mathbf{pair}(a, b)) = \Sigma(a' : A' \rho \rho' a). B'(\rho, a)(\rho', a') b$$

and $\text{fib}_{\Sigma(A, B)' } \rho \rho' (\mathbf{pair}(a, b))$ is given by closure of \mathbf{C} under dependent sum applied to $\text{fib}_{A'} \rho \rho' a$ and $\text{fib}_{B'}(\rho, a)(\rho', a') b$.

Given elements (a, a') of $(A, \langle A', \text{fib}_{A'} \rangle)$ and (b, b') of $(B, \langle B', \text{fib}_{B'} \rangle)[(a, a')]$ in \mathcal{M}^* , we define the pairing $\mathbf{pair}^*((a, a'), (b, b')) = (\mathbf{pair}(a, b), \langle a', b' \rangle)$.

Given an element $(\mathbf{pair}(a, b), \langle a', b' \rangle)$ of $\Sigma^*((A, \langle A', \text{fib}_{A'} \rangle), (B, \langle B', \text{fib}_{B'} \rangle))$ in \mathcal{M}^* , we define the projections $\mathbf{fst}^*(\mathbf{pair}(a, b), \langle a', b' \rangle) = (a, a')$ and $\mathbf{snd}^*(\mathbf{pair}(a, b), \langle a', b' \rangle) = (b, b')$.

3.2.3 Universes

We define the universe $U_n^* : \text{Type}^*(\Gamma, \Gamma')$ as $U_n^* = (U_n, \langle U_n', \text{fib}_{U_n'} \rangle)$ where $U_n' \rho \rho' A = |A| \rho \rightarrow U_n^{\text{fib}}$ and $\text{fib}_{U_n'} \rho \rho' A$ is given by $C(U_n^{\text{fib}})$ and closure of C under exponentiation (note that fibrancy of $|A| \rho$ is not used). We have carefully chosen our definitions so that we get $\text{Elem}^*((\Gamma, \Gamma'), U_n^*) = \text{Type}_n^*(\Gamma, \Gamma')$ and see that this identity is compatible with the action in \mathcal{M}^* of substitution on both sides.

3.2.4 Natural numbers

We construct a fibrant indexed inductive set \mathbb{N}' (as per Appendix B) over $|\mathbb{N}|$ (where $\mathbb{N} : \text{Type}(1)$) with constructors $0' : \mathbb{N}' 0$ and $S' : \Pi(n : |\mathbb{N}| \rho). \mathbb{N}' n \rightarrow \mathbb{N}' (S n)$. In the translation of Appendix B, the zero constructor unfolds for instance to $0'' : \text{Id}_{|\mathbb{N}|} n 0 \rightarrow \mathbb{N}' n$ for $n : |\mathbb{N}|$. This use of Id is critical to ensure fibrancy of \mathbb{N}' .

In context $(\Gamma, \Gamma') : \text{Con}^*$, we now define $\mathbb{N}^* = (\mathbb{N}, \lambda \rho \rho'. \mathbb{N}')$. We have $0^* = (0, \lambda \rho \rho'. 0')$ and $S^*(n, n') = (S(n), \lambda \rho \rho'. S' n \rho n')$ for $(n, n') : \text{Elem}((\Gamma, \Gamma'), \mathbb{N}^*)$.

Given $(P, P') : \text{Type}((\Gamma, \Gamma'). \mathbb{N}^*)$ with

$$(z, z') : \text{Elem}((\Gamma, \Gamma')(P, P')[0^*]), \quad (s, s') : \text{Elem}((\Gamma, \Gamma'). \mathbb{N}^*. (P, P')(p, S^*(q))p)$$

and $(n, n') : \text{Elem}((\Gamma, \Gamma'), \mathbb{N}^*)$, we define the elimination

$$\text{natrec}^*((P, P'), (z, z'), (s, s'), (n, n')) = (\text{natrec}(P, z, s, n), \lambda \rho \rho'. h' n \rho (n' \rho \rho'))$$

where $h' : \Pi(m : |\mathbb{N}|)(m' : \mathbb{N}' m). P'(\rho, m)(\rho', m') (\text{natrec}(P \rho^+, z \rho, s \rho^{+++}, m))$ is given by induction on \mathbb{N}' with defining equations

$$h' 0 0' = z' \rho \rho', \quad h' (S(n)) (S' n n') = s'(\rho, n, \text{natrec}(P, z, s, n))(\rho', n', h' n n').$$

3.2.5 Dependent paths

Let $\langle A, A' \rangle : \mathbb{I} \rightarrow \text{Type}^*(\Gamma, \Gamma')$ and $(a_b, a'_b) : \text{Elem}^*((\Gamma, \Gamma'), (A b, A' b))$ for $b \in \{0, 1\}$. We then define the dependent path type $\text{Path}^*(\langle A, A' \rangle, (a_0, a'_0), (a_1, a'_1)) : \text{Type}^*(\Gamma, \Gamma')$ as the tuple $(\text{Path}(A, a_0, a_1), \langle \text{Path}(A, a_0, a_1)', \text{fib}_{\text{Path}(A, a_0, a_1)'} \rangle)$ where

$$\text{Path}(A, a_0, a_1)' \rho \rho' (\langle \rangle(u)) = \text{Path}_{\lambda(i:\mathbb{I}). (A' i \rho \rho' (u i)). 1} (a'_0 \rho \rho') (a'_1 \rho \rho')$$

and $\text{fib}_{\text{Path}(A, a_0, a_1)'} \rho \rho' (\langle \rangle(u))$ is closure of C under Path applied to $(A' i \rho \rho' (u i)). 2$ for $i : \mathbb{I}$.

For $\langle u, u' \rangle : \Pi(i : \mathbb{I}). \text{Elem}^*((\Gamma, \Gamma'), (A i, A' i))$, we define path abstraction as $\langle \rangle^*(\langle u, u' \rangle) = (\langle \rangle(u), \lambda \rho \rho' i. u' i \rho \rho')$.

Given $(p, p') : \text{Elem}^*((\Gamma, \Gamma'), \text{Path}^*(\langle A, A' \rangle, (a_0, a'_0), (a_1, a'_1)))$ and $i : \mathbb{I}$, we define the path application $\text{ap}^*(p, i) = (\text{ap}(p, i), \lambda \rho \rho'. u' \rho \rho' i)$.

3.2.6 Filling operation

Given $\langle A, A' \rangle : \mathbb{I} \rightarrow \text{Type}^*(\Gamma, \Gamma')$, $\varphi : \mathbb{F}$, $b \in \{0, 1\}$, and $\langle u, u' \rangle : \Pi(i : \mathbb{I}). [\varphi] \vee (i = b) \rightarrow \text{Elem}^*((\Gamma, \Gamma'), (A i, A' i))$, we have to extend u to

$$\text{fill}^*(\langle A, A' \rangle, \varphi, b, \langle u, u' \rangle) : \Pi(i : \mathbb{I}). \text{Elem}^*((\Gamma, \Gamma'), (A i, A' i)).$$

We define $\text{fill}^*(\langle A, A' \rangle, \varphi, b, \langle u, u' \rangle) = \langle \text{fill}(A, \varphi, b, u), \text{fill}(A, \varphi, b, u)' \rangle$ where

$$\text{fill}(A, \varphi, b, u)' i \rho \rho' : A' i \rho \rho' (\text{fill}(A, \varphi, b, u) i) \rho$$

is defined using fill from (2) as

$$\text{fill}(A, \varphi, b, u)' i \rho \rho' = \text{fill}(\lambda i. A' i \rho \rho' (\text{fill}(A, \varphi, b, u) i) \rho, \varphi, b, \lambda i x. u' i x \rho \rho').$$

3.2.7 Glue types

Before defining the glueing operation in \mathcal{M}^* , we will develop several lemmas relating notions such as contractibility and equivalences in \mathcal{M} with the corresponding notions of Section 2.2. Given $f : \text{Elem}(\Gamma, A \rightarrow B)$ in \mathcal{M} , we write $|f| : \Pi(\rho : |\Gamma|). |A| \rho \rightarrow |B| \rho$ for $|f| \rho a = \mathbf{app}(f \rho, a)$. This notation overlaps with the action of $|-|$ on elements, but we will not use that one here.

Just in this subsection, we will use the alternative definition via given left and right homotopy inverses instead of contractible homotopy fibers of both equivalences \mathbf{Equiv} in the cubical cwf \mathcal{M} and equivalences \mathbf{Equiv} in the (current) internal language. In both settings, there are maps back and forth to the usual definition, which are furthermore natural in the context in the case of the cubical cwf \mathcal{M} . The statements we will prove are then also valid for the usual definition.

► **Lemma 5.** *Given $f : \text{Elem}(\Gamma, A \rightarrow B)$ in \mathcal{M} with $\text{Elem}(\Gamma, \mathbf{isEquiv}(f))$, we have $\Pi(\rho : |\Gamma|). \mathbf{isEquiv}(|f| \rho)$. This is natural in Γ .*

Proof. A (left or right) homotopy inverse $g : \text{Elem}(\Gamma, B \rightarrow A)$ to f in \mathcal{M} becomes a (left or right, respectively) homotopy inverse $|g| \rho$ to $|f| \rho$ for $\rho : |\Gamma|$. ◀

► **Lemma 6.** *Given $(f, f') : \text{Elem}((\Gamma, \Gamma'), (A, A') \rightarrow (B, B'))$ in \mathcal{M}^* , the following statements are logically equivalent, naturally in (Γ, Γ') :*

$$\text{Elem}((\Gamma, \Gamma'), \mathbf{isEquiv}^*(f, f')), \quad (3)$$

$$\text{Elem}(\Gamma, \mathbf{isEquiv}(f)) \times \Pi(\rho : |\Gamma|)(\rho' : \Gamma') \mathbf{isEquiv}(\langle |f| \rho, f' \rho \rho' \rangle), \quad (4)$$

$$\text{Elem}(\Gamma, \mathbf{isEquiv}(f)) \times \Pi(\rho : |\Gamma|)(\rho' : \Gamma') (a : |A| \rho) \mathbf{isEquiv}(f' \rho \rho' a) \quad (5)$$

where $\langle |f| \rho, f' \rho \rho' \rangle : \Sigma(a : |A| \rho) A' \rho \rho' a \rightarrow \Sigma(b : |B| \rho) B' \rho \rho' b$.

Proof. Let us only look at homotopy left inverses.

For (3) \rightarrow (4), a homotopy left inverse (g, g') to (f, f') in \mathcal{M}^* gives a homotopy left inverse $\langle |g| \rho, g' \rho \rho' \rangle$ to $\langle |f| \rho, f' \rho \rho' \rangle$ for all ρ, ρ' .

For (4) \rightarrow (5), we use Lemma 5 and note that a fiberwise map over an equivalence is a fiberwise equivalence exactly if it is an equivalence on total spaces (the corresponding statement for identity types instead of paths is [27, Theorem 4.7.7]).

For (5) \rightarrow (3), given a homotopy left inverse g to the equivalence f in \mathcal{M} and a homotopy left inverse $\bar{g}' \rho \rho' a : B' \rho \rho' (|f| a) \rightarrow A' \rho \rho' a$ to $f' \rho \rho' a$ for all ρ, ρ', a , we use Lemma 5 to transpose \bar{g}' to the second component $g' \rho \rho' b : B' \rho \rho' b \rightarrow A' \rho \rho' (|g| b)$ for all ρ, ρ', b of a homotopy left inverse (g, g') to (f, f') in \mathcal{M}^* . ◀

We can now define glue types in \mathcal{M}^* . Let $(A, A') : \text{Type}(\Gamma, \Gamma')$, $\varphi : \mathbb{F}$, $b \in \{0, 1\}$, $\langle T, T' \rangle : [\varphi] \rightarrow \text{Type}(\Gamma, \Gamma')$, and $\langle e, e' \rangle : [\varphi] \rightarrow \text{Elem}((\Gamma, \Gamma'), \mathbf{Equiv}^*((T \text{tt}, T' \text{tt}), (A, A')))$.

We define $\mathbf{Glue}^*((A, A'), \varphi, b, \langle T, T' \rangle, \langle e, e' \rangle) = (\mathbf{Glue}(A, \varphi, b, T, e), \langle G', \mathbf{fib}_{G'} \rangle)$ where

$$G' \rho \rho' (\mathbf{glue}(a, t)) = \mathbf{Glue}(A' \rho \rho' a).1 [\varphi \mapsto (T' \text{tt} \rho \rho' (t \text{tt}), ((e' \text{tt} \rho \rho').1, w \text{tt} \rho \rho'))]$$

using the witness $w \text{tt} \rho \rho'$ that $(e' \text{tt} \rho \rho').1$ is an equivalence provided by the direction from (3) to (5) of Lemma 6 and $\mathbf{fib}_{G'} \rho \rho' (\mathbf{glue}(a, t))$ is given by closure of \mathbf{C} under \mathbf{Glue} applied to $(A' \rho \rho' a).2$ and $T' \text{tt} \rho, \rho' (t \text{tt})$ on $[\varphi]$.

We define $\mathbf{unglue}^* = (\mathbf{unglue}, \mathbf{unglue}')$ where $\mathbf{unglue}' \rho \rho' (\mathbf{glue}(a, t)) = \mathbf{unglue}$.

Given $(a, a') : \text{Elem}((\Gamma, \Gamma'), (A, A'))$ and $(t, t') : [\varphi] \rightarrow \text{Elem}((\Gamma, \Gamma'), (T \text{tt}, T' \text{tt}))$ such that $\mathbf{app}^*(\mathbf{fst}^*(e, e') \text{tt}, (t, t') \text{tt}) = (a, a')$ on $[\varphi]$, we define $\mathbf{glue}^*((a, a'), (t, t'))$ as the pair $(\mathbf{glue}(a, t), \mathbf{glue}(a, t'))$ where $\mathbf{glue}(a, t') \rho \rho' = \mathbf{glue}(a' \rho \rho') [\varphi \mapsto t' \text{tt} \rho \rho']$.

3.3 Main result

One checks in a mechanical fashion that the operations we have defined above satisfy the required laws, including stability under substitution in the context (Γ, Γ') . We thus obtain the following statement.

► **Theorem 7** (Scoring). *Assume the parameters $\mathcal{C}, \mathbb{I}, \mathbb{F}$ satisfy the assumptions of Section 2.2. Then given any cubical cwf \mathcal{M} , the scoring \mathcal{M}^* is a cubical cwf with operations defined as above. We further have morphism $\mathcal{M}^* \rightarrow \mathcal{M}$ of cubical cwf given by the first projection.*

4 Homotopy canonicity

► **Theorem 8** (Homotopy canonicity). *Assume the parameters $\mathcal{C}, \mathbb{I}, \mathbb{F}$ satisfy the assumptions of Section 2.2. In the internal language of presheaves over \mathcal{C} , given a closed natural $n : \text{Elem}(1, \mathbb{N})$ in the term model \mathcal{T} , we have a numeral $k : \mathbb{N}$ with a path term $p : \text{Elem}(1, \text{Path}(\mathbb{N}, n, \mathbb{S}^k(0)))$.*

Proof. We start the arguing reasoning externally. Using Theorem 7, we build the scoring \mathcal{T}^* of \mathcal{T} . Since \mathcal{T} is the initial cubical cwf by Theorem 2, we obtain a section F of the cubical cwf morphism $\mathcal{T}^* \rightarrow \mathcal{T}$.

Let us now proceed in the internal language. Recall the construction of Section 3.2.4 of natural numbers in \mathcal{T}^* . We observe that $\Sigma(n : |\mathbb{N}|). \mathbb{N}' n$ forms a natural number type (in the terminal context) in \mathcal{S} . It is thus homotopy equivalent to \mathbb{N} . Under this equivalence, the first projection $\Sigma(n : |\mathbb{N}|). \mathbb{N}' n \rightarrow |\mathbb{N}|$ implements the map sending $k : \mathbb{N}$ to $\mathbb{S}^k(0)$.

Inspecting the action of F on $n : \text{Elem}(1, \mathbb{N})$, we obtain $n' : \mathbb{N}', n$. By the preceding paragraph, this corresponds to $k : \mathbb{N}$ with a path $p' : \mathbb{I} \rightarrow |\mathbb{N}|$ from n to $\mathbb{S}^k(0)$. Now $p = \langle \rangle(p')$ is the desired witness of homotopy canonicity. ◀

5 Extensions

5.1 Identity types

Our treatment extends to the variation of cubical cwf that includes identity types.

Identity types in a cubical cwf denoted as in Section 1.3 consist of the following operations and laws (omitting stability under substitution), internal to presheaves over \mathcal{C} . Fix A in $\text{Type}(\Gamma)$. Given x, y in $\text{Elem}(\Gamma, A)$, we have $\text{Id}(A, x, y)$ in $\text{Type}(\Gamma)$, of level n if A is. Given a in $\text{Elem}(\Gamma, A)$, we have $\text{refl}(a)$ in $\text{Elem}(\Gamma, \text{Id}(A, a, a))$. Given P in $\text{Type}(\Gamma.A.\text{Ap}.\text{Id}(\text{App}, \text{qp}, \text{q}))$ and d in $\text{Elem}(\Gamma.A, P[\text{q}, \text{q}, \text{refl}(q)])$ and x, y in $\text{Elem}(\Gamma, A)$ and p in $\text{Elem}(\Gamma, \text{Id}(A, x, y))$, we have $J(P, d, x, y, p)$ in $\text{Elem}(\Gamma, P[x, y, p])$. We have $J(P, d, a, a, \text{refl}(a)) = d[a]$.

We can interpret univalent type theory in such any such cubical cwf as per Remark 1.

The standard model of Section 2.3 has identity type $\text{Id}(\langle A, \text{fib}_A \rangle, x, y) : \text{Type}(\Gamma)$ given by $\Pi(\rho : \Gamma).\text{Id}_{A\rho}(x\rho)(y\rho)$ using Andrew Swan's construction of Id referenced in Appendix B. We omit the evident description of the remaining operations.

To obtain homotopy canonicity in this setting, it suffices to extend the scoring construction \mathcal{M}^* of Section 3 to identity types. Given $A : \text{Type}(1)$ and $A' : |A| \rightarrow \mathbb{U}_\omega^{\text{fib}}$, we define $\text{Id}'_{A, A'}$ as the fibrant indexed inductive set (as per Appendix B) over $x, y : |A|, p : |\text{Id}(A, x, y)|, x' : A' x, y' : A' y$ with constructor $\text{refl}' : \Pi(a : |A|)(a' : A' a).\text{Id}'_{A, A'} a a (\text{refl}(a)) a' a'$.

Now fix $(A, A') : \text{Type}^*(\Gamma, \Gamma')$. Given $\rho : |\Gamma|$ and $\rho' : \Gamma' \rho$, Given elements $(x, x'), (y, y')$ of (A, A') in \mathcal{M}^* , we define

$$\text{Id}^*((A, A'), (x, x'), (y, y')) = (\text{Id}(A, x, y), \lambda \rho \rho' p.\text{Id}'_{A\rho, A'\rho\rho'} x\rho y\rho p (x' \rho \rho') (y' \rho \rho')).$$

Given an element (a, a') of (A, A') in \mathcal{M}^* , we define $\mathbf{refl}^*(a, a') = (\mathbf{refl}(a), \mathbf{refl}(a'))$ where $\mathbf{refl}(a)' \rho \rho' = \mathbf{refl}' a \rho (a' \rho \rho')$. The eliminator $\mathbf{J}((C, C'), (d, d'), (x, x'), (y, y'), (p, p'))$ is defined as $(\mathbf{J}(C, d, x, y, p), \lambda \rho \rho'. h' x \rho y \rho p \rho (x' \rho \rho') (y' \rho \rho') (p' \rho \rho'))$ where

$$h' : \Pi(x y : |A| \rho)(p : |\mathbf{Id}(A, x, y)| \rho)(x' : A' \rho')(y' : A' \rho')(p' : \mathbf{Id}'_{A \rho, A' \rho \rho'} x y p x' y').$$

$$P'(\rho, x, y, p)(\rho', x', y, p')(\mathbf{J}(P \rho^{+++}, d \rho^+, x, y, p))$$

is given by induction on $\mathbf{Id}'_{A \rho, A' \rho \rho'}$ via $h' a a (\mathbf{refl}(a)) a' a' (\mathbf{refl}' a a') = d'(\rho, a)(\rho', a')$.

5.2 Higher inductive types

Our treatment extends to higher inductive types [27], following the semantics presented in [5]. Crucially, we have fibrant *indexed* higher inductive sets in presheaves over \mathcal{C} as we have what we would call fibrant *uniformly indexed* higher inductive sets in the same fashion as in [5] and fibrant identity sets [4, 20], mirroring the derivation of fibrant indexed inductive sets from fibrant uniformly indexed inductive sets and fibrant identity sets recollected in Appendix B.⁵

Let us look at the case of the *suspension* operation in a cubical cwf, where $\mathbf{Susp}(A) : \mathbf{Type}(\Gamma)$ has constructors \mathbf{north} , \mathbf{south} and $\mathbf{merid}(a, i)$ for $a : A$ and $i : \mathbb{I}$ with $\mathbf{merid}(a, 0) = \mathbf{north}$ and $\mathbf{merid}(a, 1) = \mathbf{south}$.

For the scoping model of Section 3, we define for $A : \mathbf{Type}(1)$ and $A' : |A| \rightarrow \mathbf{U}_\omega^{\text{fib}}$ the indexed higher inductive set $\mathbf{Susp}'_{A, A'}$ over $|\mathbf{Susp}(A)|$ with constructors

$$\mathbf{north}' : \mathbf{Susp}'_{A, A'} \mathbf{north} \qquad \mathbf{south}' : \mathbf{Susp}'_{A, A'} \mathbf{south}$$

$$\mathbf{merid}' a a' i : (\mathbf{Susp} A)'(\mathbf{merid}(a, i))[i = 0 \mapsto \mathbf{north}', i = 1 \mapsto \mathbf{south}']$$

for $a : |a|$ and $a' : A' a$ and $i : \mathbb{I}$ (using the notation of [5]). In the above translation to a uniformly indexed higher inductive set, the constructor \mathbf{north}' will for example be replaced by $\mathbf{north}'' : \mathbf{Id}_{|\mathbf{Susp}(A)|} u \mathbf{north} \rightarrow \mathbf{Susp}'_{A, A'} u$.

Given $(A, A') : \mathbf{Type}(\Gamma, \Gamma')$, we then define $\mathbf{Susp}^*(A, A') = (\mathbf{Susp}(A), \lambda \rho \rho'. \mathbf{Susp}'_{A \rho, A' \rho \rho'})$, with constructors and eliminator treated as in Section 5.1.

References

- 1 P. Aczel. On Relating Type Theories and Set Theories. In T. Altenkirch, B. Reus, and W. Naraschewski, editors, *Types for Proofs and Programs*, volume 1657 of *Lecture Notes in Computer Science*, pages 1–18. Springer Verlag, Berlin, Heidelberg, New York, 1999.
- 2 C. Angiuli, G. Brunerie, T. Coquand, K.-B. Hou (Favonia), B. Harper, and D. Licata. Cartesian cubical type theory. preprint available from the home page of D. Licata, 2017.
- 3 M. Bezem, T. Coquand, and S. Huber. A Model of Type Theory in Cubical Sets. In R. Matthes and A. Schubert, editors, *19th International Conference on Types for Proofs and Programs (TYPES 2013)*, volume 26 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 107–128. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2014.
- 4 C. Cohen, T. Coquand, S. Huber, and A. Mörtberg. Cubical Type Theory: A Constructive Interpretation of the Univalence Axiom. In T. Uustalu, editor, *21st International Conference on Types for Proofs and Programs (TYPES 2015)*, volume 69 of *Leibniz International Proceedings in Informatics (LIPIcs)*. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2018.

⁵ We stress that the use of “set” in this context refers to the types of the language of presheaves over \mathcal{C} , not homotopy sets.

- 5 T. Coquand, S. Huber, and A. Mörtberg. On higher inductive types in cubical type theory. In *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '18*, pages 255–264, New York, NY, USA, 2018. ACM.
- 6 P. Dybjer. Internal Type Theory. In *Lecture Notes in Computer Science*, pages 120–134. Springer Verlag, Berlin, Heidelberg, New York, 1996.
- 7 S. Eilenberg. On the relation between the fundamental group of a space and the higher homotopy groups. *Fundamenta Mathematicae*, 32(1):169–175, 1939.
- 8 P. Gabriel and M. Zisman. *Calculus of fractions and homotopy theory*, volume 35 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, 1967.
- 9 N. Gambino and J. Kock. Polynomial functors and polynomial monads. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 154, pages 153–192. Cambridge University Press, 2013.
- 10 N. Gambino and C. Sattler. The Frobenius condition, right properness, and uniform fibrations. *Journal of Pure and Applied Algebra*, 221(12):3027–3068, 2017.
- 11 K. Gödel. Über eine bisher noch nicht benützte Erweiterung des finiten Standpunkts. *Dialectica*, 12:280–287, 1958.
- 12 M. Hofmann. Syntax and semantics of dependent types. In A.M. Pitts and P. Dybjer, editors, *Semantics and logics of computation*, volume 14 of *Publ. Newton Inst.*, pages 79–130. Cambridge University Press, Cambridge, 1997.
- 13 S. Huber. Canonicity for cubical type theory. *Journal of Automated Reasoning*, 2018.
- 14 A. Joyal. Notes on clans and tribes, 2017.
- 15 C. Kapulkin and P. LeFanu Lumsdaine. The simplicial model of univalent foundations (after Voevodsky). *arXiv preprint arXiv:1211.2851*, 2012.
- 16 K. Kapulkin and V. Voevodsky. Cubical approach to straightening. preprint available from the home page of K. Kapulkin, 2018.
- 17 G.M. Kelly. A unified treatment of transfinite constructions for free algebras, free monoids, colimits, associated sheaves, and so on. *Bulletin of the Australian Mathematical Society*, 22(1):1–83, 1980.
- 18 D. R. Licata, I. Orton, A. M. Pitts, and B. Spitters. Internal Universes in Models of Homotopy Type Theory. In H el ene Kirchner, editor, *3rd International Conference on Formal Structures for Computation and Deduction (FSCD 2018)*, volume 108 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 22:1–22:17. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2018.
- 19 P. LeFanu Lumsdaine and M. Shulman. Semantics of higher inductive types. *arXiv preprint arXiv:1705.07088*, 2017.
- 20 I. Orton and A. M. Pitts. Axioms for modelling cubical type theory in a topos. In *25th EACSL Annual Conference on Computer Science Logic (CSL 2016)*, volume 62 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 24:1–24:19, Dagstuhl, Germany, 2016. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
- 21 E. Riehl and M. Shulman. A type theory for synthetic ∞ -categories. *Higher Structures*, 1(1), 2018.
- 22 M. Shulman. Univalence for inverse diagrams and homotopy canonicity. *Mathematical Structures in Computer Science*, 25(5):1203–1277, 2015.
- 23 T. Streicher. *Semantics of Type Theory*. Progress in Theoretical Computer Science. Birkh user Basel, 1991.
- 24 A. Swan. An algebraic weak factorisation system on 01-substitution sets: A constructive proof. *Journal of Logic & Analysis*, 8(1):1–35, 2016.
- 25 A. Swan. Semantics of higher inductive types, 2017. On the HoTT mailing list.
- 26 W. W. Tait. Intensional interpretations of functionals of finite type I. *Journal of Symbolic Logic*, 32(2):198–212, 1967.
- 27 The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. <http://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.

28 V. Voevodsky. The equivalence axiom and univalent models of type theory. (Talk at CMU on February 4, 2010). Preprint arXiv:1402.5556 [math.LO], 2014.

A Rules of the term model

We denote the objects of our base category \mathcal{C} by X, Y, Z and its morphisms by f, g, h . In the term model \mathcal{T} morphisms $f: Y \rightarrow X$ act on judgments at stage X via an implicit substitution, while for substitutions on object variables we will use explicit substitutions. For this to make sense we first define the raw expressions as a presheaf: at stage X this is given by

$$\begin{aligned}
 \Gamma, \Delta &::= \varepsilon \mid \Gamma.A \\
 A, B, t, u, v &::= \mathbf{q} \mid t\sigma \mid \mathbf{U}_n \mid \Pi(A, B) \mid \mathbf{lam}(u) \mid \mathbf{app}(u, v) \\
 &\mid \Sigma(A, B) \mid \mathbf{pair}(u, v) \mid \mathbf{fst}(u) \mid \mathbf{snd}(u) \\
 &\mid \mathbf{Path}(\bar{A}, u, v) \mid \langle \bar{u} \mid \mathbf{ap}(u, r) \\
 &\mid \mathbf{Glue}(A, \varphi, \bar{B}, \bar{u}) \mid \mathbf{glue}(v, \bar{u}) \mid \mathbf{unglue}(u) \\
 &\mid \mathbf{fill}(\bar{A}, \varphi, b, \bar{u}, r) \mid \dots \\
 \bar{A}, \bar{B}, \bar{u}, \bar{v} &::= (A_{f,r})_{f,r} \mid (A_f)_{f \in [\varphi]} \\
 \sigma, \tau, \delta &::= \mathbf{p} \mid \mathbf{id} \mid \sigma\tau \mid (\sigma, u) \mid ()
 \end{aligned}$$

where $b \in \{0, 1\}$, $\varphi \in \mathbb{F}(X)$, and we skipped the constants for natural numbers. Above, we have families of expressions, say $\bar{A} = (A_{f,r})_{f,r}$, whose index set ranges over certain Y , $f: Y \rightarrow X$, and $r \in \mathbb{I}(Y)$, and $A_{f,r}$ is a raw expression at stage Y ; likewise $(A_f)_{f \in [\varphi](X)}$ consists of raw expressions A_f at stage Y for $f: Y \rightarrow X$ in the sieve $[\varphi]$ on X . (The exact index sets will be clear from the typing rules below.) All other occurrences of r above have $r \in \mathbb{I}(X)$. The restrictions along $f: Y \rightarrow X$ on the raw syntax then leave all the usual cwf structure untouched, so we have $\mathbf{q}f = \mathbf{q}$ and $(\Pi(A, B))f = \Pi(Af, Bf)$, and uses the restrictions in \mathbb{I} and \mathbb{F} accordingly, e.g., $(\mathbf{ap}(u, r))f = \mathbf{ap}(uf, rf)$, and we will re-index families according to $\bar{A}f = (A_{gf,rf})$ for $\bar{A} = (A_{g,r})_{g,r}$.

To get the *initial* cubical cwf we in fact need more annotations to the syntax in order to be able to define a partial interpretation (cf. [23, 12]) on the raw syntax. But to enhance readability we suppress these annotations.

We will now describe a type system indexed by stages X . The forms of judgment are:

$$\Gamma \vdash_X \quad \Gamma \vdash_X A \quad \Gamma \vdash_X A = B \quad \Gamma \vdash_X t : A \quad \Gamma \vdash_X t = u : A \quad \sigma : \Delta \rightarrow_X \Gamma$$

where the involved expressions are at stage X .

► **Remark 9.** In cubical type theory as described in [4] we did not index judgments by objects X but allowed extending context by interval variables instead. Loosely speaking, a judgment $\Gamma \vdash_{\{i_1, \dots, i_n\}} \mathcal{J}$ corresponds to $i_1 : \mathbb{I}, \dots, i_n : \mathbb{I}, \Gamma \vdash \mathcal{J}$ given the setting of [4].

As mentioned above we have the rule:

$$\frac{\Gamma \vdash_X \mathcal{J} \quad f: Y \rightarrow X}{\Gamma f \vdash_Y \mathcal{J}f}$$

At each stage we have all the usual rules valid in a cwf with Π -types, Σ -types, universes, and natural numbers. We will present some of the rules, but skip all congruence rules.

$$\frac{}{\varepsilon \vdash_X} \quad \frac{\Gamma \vdash_X \quad \Gamma \vdash_X A}{\Gamma.A \vdash_X} \quad \frac{\Gamma \vdash_X A \quad \sigma : \Delta \rightarrow_X \Gamma}{\Delta \vdash_X A\sigma} \quad \frac{\Gamma \vdash_X t : A \quad \sigma : \Delta \rightarrow_X \Gamma}{\Delta \vdash_X t\sigma : A\sigma}$$

$$\frac{\Gamma \vdash_X A}{\Gamma.A \vdash_X \mathbf{q} : \mathbf{Ap}} \quad \frac{\Gamma \vdash_X t : A \quad \Gamma \vdash_X A = B}{\Gamma \vdash_X t : B} \quad \frac{\Gamma \vdash_X}{\mathbf{id} : \Gamma \rightarrow_X \Gamma} \quad \frac{\Gamma \vdash_X}{(): \Gamma \rightarrow \varepsilon}$$

$$\begin{array}{c}
\frac{\Gamma \vdash_X A}{\mathbf{p}: \Gamma.A \rightarrow_X \Gamma} \qquad \frac{\sigma: \Delta \rightarrow_X \Gamma \quad \tau: \Theta \rightarrow_X \Delta}{\sigma\tau: \Theta \rightarrow_X \Gamma} \\
\\
\frac{\sigma: \Delta \rightarrow_X \Gamma \quad \Gamma \vdash_X A \quad \Delta \vdash_X u: A\sigma}{(\sigma, u): \Delta \rightarrow_X \Gamma.A} \\
\\
\frac{\Gamma.A \vdash_X B}{\Gamma \vdash_X \Pi(A, B)} \quad \frac{\Gamma.A \vdash_X B \quad \Gamma.A \vdash_X b: B}{\Gamma \vdash_X \mathbf{lam}(b): \Pi(A, B)} \quad \frac{\Gamma \vdash_X w: \Pi(A, B) \quad \Gamma \vdash_X u: A}{\Gamma \vdash_X \mathbf{app}(w, u): B[u]}
\end{array}$$

where we write $[u]$ for (id, u) and σ^+ for $(\sigma\mathbf{p}, \mathbf{q})$. The judgmental equalities (skipping suitable premises, types, and contexts) are:

$$\begin{array}{l}
\text{id } \sigma = \sigma \text{ id} = \sigma \quad (\sigma\tau)\delta = \sigma(\tau\delta) \quad ()\sigma = () \quad (\sigma, u)\delta = (\sigma\delta, u\delta) \quad \mathbf{p}(\sigma, u) = \sigma \\
\mathbf{q}(\sigma, u) = u \quad (\mathbf{p}, \mathbf{q}) = \text{id} \quad A \text{ id} = A \quad (A\sigma)\delta = A(\sigma\delta) \quad u \text{ id} = u \quad (u\sigma)\delta = u(\sigma\delta) \\
(\Pi(A, B))\sigma = \Pi(A\sigma, B\sigma^+) \quad (\mathbf{lam}(b))\sigma = \mathbf{lam}(b\sigma^+) \quad \mathbf{app}(w, u)\delta = \mathbf{app}(w\delta, u\delta) \\
\mathbf{app}(\mathbf{lam}(b), u) = b[u] \quad w = \mathbf{lam}(\mathbf{app}(w\mathbf{p}, \mathbf{q}))
\end{array}$$

We skip the rules for Σ -types and natural numbers as they are standard, but simply indexed with an object X as we did for Π -types. The rules for universes are:

$$\frac{\Gamma \vdash_X}{\Gamma \vdash_X \mathbb{U}_n} \quad \frac{\Gamma \vdash_X}{\Gamma \vdash_X \mathbb{U}_n : \mathbb{U}_{n+1}} \quad \frac{\Gamma \vdash_X A : \mathbb{U}_n}{\Gamma \vdash_X A : \mathbb{U}_{n+1}} \quad \frac{\Gamma \vdash_X A : \mathbb{U}_n}{\Gamma \vdash_X A}$$

and we skip the rules for equality and closure under the type formers Π, Σ , natural numbers, \mathbf{Path} , and \mathbf{Glue} .

To state the rules for dependent path-types we introduce the following abbreviations. We write $\Gamma.\mathbb{I} \vdash_X \bar{A}$ if $\bar{A} = (A_{f,r})$ is a family indexed by Y , $f: Y \rightarrow X$, and $r \in \mathbb{I}(Y)$ such that

$$\Gamma f \vdash_Y A_{f,r} \text{ and } \Gamma f g \vdash_Z (A_{f,r})g = A_{fg,r}.$$

Given $\Gamma.\mathbb{I} \vdash_X \bar{A}$ we write $\Gamma.\mathbb{I} \vdash_X \bar{u} : \bar{A}$ whenever $\bar{u} = (u_{f,r})$ is a family indexed by Y , $f: Y \rightarrow X$, and $r \in \mathbb{I}(Y)$ such that

$$\Gamma f \vdash_Y u_{f,r} : A_{f,r} \text{ and } \Gamma f g \vdash_Z (u_{f,r})g = u_{fg,rg} : A_{fg,rg}.$$

The rules for the dependent path type are:

$$\frac{\Gamma.\mathbb{I} \vdash_X \bar{A} \quad \Gamma \vdash_X u : A_{\text{id}_X,0} \quad \Gamma \vdash_X v : A_{\text{id}_X,1}}{\Gamma \vdash_X \mathbf{Path}(\bar{A}, u, v)} \quad \frac{\Gamma.\mathbb{I} \vdash_X \bar{A} \quad \Gamma.\mathbb{I} \vdash_X \bar{u} : \bar{A}}{\Gamma \vdash_X \mathbf{lam}(\bar{u}) : \mathbf{Path}(\bar{A}, u_{\text{id}_X,0}, u_{\text{id}_X,1})} \\
\frac{\Gamma \vdash_X t : \mathbf{Path}(\bar{A}, u, v) \quad r \in \mathbb{I}(X)}{\Gamma \vdash_X \mathbf{ap}(t, r) : A_{\text{id}_X,r}}$$

$$\begin{array}{l}
\mathbf{ap}(\mathbf{lam}(\bar{u}), r) = u_{\text{id},r} \quad t = \mathbf{lam}(\mathbf{ap}(tf, r)_{f,r}) \quad \mathbf{Path}(\bar{A}, u, v)\sigma = \mathbf{Path}((A_{f,r}\sigma f)_{f,r}, u\sigma, v\sigma) \\
(\mathbf{lam}(\bar{u}))\sigma = \mathbf{lam}((u_{f,r}\sigma f)_{f,r}) \quad (\mathbf{ap}(t, r))\sigma = \mathbf{ap}(t\sigma, r)
\end{array}$$

Note that in general these rules might have infinitely many premises. We get the non-dependent path type for $\Gamma \vdash_X A$ by using the family $A_{f,r} := Af$.

23:18 Homotopy canonicity for cubical type theory

Given $\Gamma.\mathbb{I} \vdash_X \bar{A}$ and $b \in \{0, 1\}$ we write $\Gamma.\mathbb{I} \vdash_X^{\varphi, b} \bar{u} : \bar{A}$ for $\bar{u} = (u_{f,r})$ a family indexed over all $Y, f: Y \rightarrow X$, and $r \in \mathbb{I}(Y)$ such that either f is in the sieve $[\varphi]$ or $r = b$ and we have

$$\Gamma f \vdash_Y u_{f,r} : A_{f,r} \text{ and } \Gamma f g \vdash_Z (u_{f,r})g = u_{fg,rg} : A_{fg,rg}$$

for all $g: Z \rightarrow Y$. The rule for the filling operation is given by:

$$\frac{\Gamma.\mathbb{I} \vdash_X \bar{A} \quad \varphi \in \mathbb{F}(X) \quad b \in \{0, 1\} \quad \Gamma.\mathbb{I} \vdash_X^{\varphi, b} \bar{u} : \bar{A} \quad r \in \mathbb{I}(X)}{\Gamma \vdash_Y \mathbf{fill}(\bar{A}, \varphi, b, \bar{u}, r) : A_{\text{id},r}}$$

with judgmental equality

$$\mathbf{fill}(\bar{A}, \varphi, b, \bar{u}, r) = u_{\text{id},r} \text{ whenever } [\varphi] \text{ is the maximal sieve or } r = b.$$

For the glueing operation we only present the formation rule; the other rules are similar as in [4] but adapted to our setting. We write $\Gamma \vdash_X^\varphi \bar{B}$ if \bar{B} is a family of B_f for $f: Y \rightarrow X$ in $[\varphi]$ with $\Gamma f \vdash_Y B_f$ which is compatible, i.e. $\Gamma f g \vdash_Z B_f g = B_{fg}$. In this case, we write likewise $\Gamma \vdash_X \bar{u} : \bar{B}$ if \bar{u} is a compatible family of terms $\Gamma f \vdash_Y u_f : B_f$.

$$\frac{\Gamma \vdash_X A \quad \varphi \in \mathbb{F}(X) \quad \Gamma \vdash_X^\varphi \bar{B} \quad \Gamma \vdash_X^\varphi \bar{u} : \mathbf{isEquiv}(\bar{B}, A)}{\Gamma \vdash_X \mathbf{Glue}(A, \varphi, \bar{B}, \bar{u})}$$

and the judgmental equality $\mathbf{Glue}(A, \varphi, \bar{B}, \bar{u}) = B_{\text{id}}$ in case $[\varphi]$ is the maximal sieve, and an equation for substitution.

This formal system gives rise to a cubical cwf \mathcal{T} as follows. First, define judgmental equality for contexts and substitutions as usual (we could also have those as primitive judgments). Next, we define presheaves \mathbf{Con} and \mathbf{Hom} on \mathcal{C} by taking, say, $\mathbf{Con}(X)$ equivalence classes $[\Gamma]_\sim$ of Γ with $\Gamma \vdash_X$ modulo judgmental equality; restrictions are induced by the (implicit) substitution: $[\Gamma]_\sim f = [\Gamma f]_\sim$. Types $\mathbf{Type}(X, [\Gamma]_\sim)$ are equivalence classes of A with $\Gamma \vdash_X A$ modulo judgmental equality, and elements are defined similarly as equivalence classes.

For type formers in \mathcal{T} let us look at path types: we have to give an element of $\mathbf{Type}(\Gamma)$ in a context (w.r.t. the internal language) $\Gamma : \mathbf{Con}, A : \mathbb{I} \rightarrow \mathbf{Type}(\Gamma), u : \mathbf{Elem}(\Gamma, A 0), v : \mathbf{Elem}(\Gamma, A 1)$. Unfolding the use of internal language, given $[\Gamma]_\sim \in \mathbf{Con}(X)$, a compatible family $[A_{f,r}]_\sim \in \mathbf{Type}(Y, [\Gamma]_\sim f)$ (for $f: Y \rightarrow X$ and $r \in \mathbb{I}(Y)$) and elements $[u]_\sim \in \mathbf{Elem}([\Gamma]_\sim, [A_{\text{id},0}]_\sim)$ and $[v]_\sim \in ([\Gamma]_\sim, [A_{\text{id},1}]_\sim)$, we have to give an element of $\mathbf{Type}(X, [\Gamma]_\sim)$, which we do by the formation rule for \mathbf{Path} .

The remainder of the cubical cwf structure for \mathcal{T} is defined in a similar manner, in fact the rules are designed to reflect the laws of cubical cwfs, and we can follow a similar argument as in [23] to show that \mathcal{T} is the *initial* cubical cwf. Given a cubical cwf \mathcal{M} over $\mathcal{C}, \mathbb{I}, \mathbb{F}$ we first have to define *partial* interpretations of the raw syntax and then show that each derivable judgment has a defined interpretation in \mathcal{M} , and for equality judgments both sides of the equation have a defined interpretation in \mathcal{M} and are equal. The partial interpretation $\llbracket - \rrbracket$ assigns meanings to raw judgments with the following signature:

$$\begin{aligned} \llbracket \Gamma \vdash_X \rrbracket &\in \mathbf{Con}_{\mathcal{M}}(X) \\ \llbracket \sigma : \Delta \rightarrow \Gamma \rrbracket &\in \mathbf{Hom}_{\mathcal{M}}(X, \llbracket \Delta \vdash_X \rrbracket, \llbracket \Gamma \vdash_X \rrbracket) \\ \llbracket \Gamma \vdash_X A \rrbracket &\in \mathbf{Type}_{\mathcal{M}}(X, \llbracket \Gamma \vdash_X \rrbracket) \\ \llbracket \Gamma \vdash_X u : A \rrbracket &\in \mathbf{Elem}_{\mathcal{M}}(X, \llbracket \Gamma \vdash_X \rrbracket, \llbracket \Gamma \vdash_X A \rrbracket) \end{aligned}$$

where among the conditions for the interpretation on the left-hand side to be defined is that all references to the interpretation on the right-hand side are defined. This proceeds by

structural induction on the raw syntax and for $\llbracket \Gamma \vdash_X \mathcal{J} \rrbracket$ to be defined we assume all the ingredients needed are already defined. E.g. for the path type $\llbracket \Gamma \vdash_X \text{Path}(\bar{A}, u, v) \rrbracket$ we in particular have to assume that the assignment $f, r \mapsto \llbracket \Gamma f \vdash_Y A_{f,r} \rrbracket$ is defined and gives rise to a suitable input of $\text{Path}_{\mathcal{M}}$.

B Indexed inductive sets in presheaves over \mathcal{C}

We work in the setting of Section 2.2 given by presheaves over \mathcal{C} .

Given a set I , a family A over I , a family B over $i : I$ and $a : A i$, and a map

$$s : \Pi(i : I)(a : A i).B i a \rightarrow I,$$

the *indexed inductive set* $W_{I,A,B,s}$ is the initial algebra of the polynomial endofunctor [9] on the (internal) category of families over I sending a family X to the family

$$\llbracket I, A, B, s \rrbracket i = \Sigma(a : A i). \Pi(b : B i a). X(s i a b).$$

Its constructive justification as an operation in the internal language of the presheaf topos using inductive constructions of the metatheory is folklore (in a classical setting, one would use transfinite colimits [17]). If I, A, B are small with respect to a universe U_i with $i \in \{0, 1, \dots, \omega\}$, then $W_{I,A,B,s} : I \rightarrow U_i$.

Let I, A, B now be small with respect to U_ω . Given elements of $\text{Fill}(I, A)$ and $\text{Fill}(\Sigma(i : I).A i, \lambda(i, a).B i a)$, we may use induction (i.e. the universal property of $W_{I,A,B,s}$) to derive an element of $\text{Fill}(I, W_{I,A,B,s})$. As in Section 2.2 for dependent products, this implies (using external reasoning) the internal statement $C(\text{Fill}(W_{I,A,B,s}))$ given $C(A i)$ for $i : I$ and $C(B i a)$ for additionally $a : I i$. We then call $W(I, A, B, s)$ a *fibrant uniformly indexed inductive set*. The qualifier *uniformly indexed* indicates that A is a fibrant family over I rather than a fibrant set with a “target” map to I that indicates the target sort of the constructor sup .

Given $A : U_\omega$ with $\text{Fill}(1, A)$, we may use the technique of Andrew Swan [24, 20] to construct a (level preserving) identity set $\text{Id}_A a_0 a_1$ for $a_0, a_1 : A$ (different from the equality set $a_0 = a_1$) with $\text{Fill}(A \times A, \lambda(a_0, a_1).\text{Id}_A a_0 a_1)$ that has the usual elimination with respect to families $C : \Pi(a_0 a_1 : A).\text{Id}_A a_0 a_1 \rightarrow U_\omega$ that satisfy $\text{Fill}(\Sigma(a_0 a_1 : A).\text{Id}_A a_0 a_1, C)$. Using external reasoning as before, one has $C(\text{Id}_A a_0 a_1)$ given $C(A)$, i.e. justifying calling $\text{Id}_A a_0 a_1$ a *fibrant identity set*; using (2) one has elimination with respect to families C of the previous signature with $C(C a_0 a_1 p)$ for all a_0, a_1, p .

Using a folklore technique, we may use fibrant identity sets to derive *fibrant indexed inductive sets* from fibrant uniformly indexed inductive sets, by which we mean the following. Given $(I, \text{fib}_I) : U_\omega^{\text{fib}}$, $(A, \text{fib}_A) : U_\omega^{\text{fib}}$, $\langle B, \text{fib}_B \rangle : A \rightarrow U_\omega^{\text{fib}}$ with maps $t : A \rightarrow I$ and $s : \Pi(a : A).B a \rightarrow I$, we have $\langle W, \text{fib}_W \rangle : I \rightarrow U_\omega^{\text{fib}}$ with

$$\text{sup} : \Pi(a : A)(f : \Pi(b : B a).W(s a b)).W(t a).$$

Given $\langle P, \text{fib}_P \rangle : \Pi(i : I).W \rightarrow U_\omega^{\text{fib}}$ with

$$h : \Pi(a : A)(f : \Pi(b : B a).W(s a b)).(\Pi(b : B a).P(s a b)(f b)) \rightarrow P(t a)(\text{sup } a f),$$

we have $v : \Pi(i : I)(w : W i).P i w$ such that

$$v(t a)(\text{sup } a f) = h a f(\lambda b.v(s a b)(f b)).$$

We will use these fibrant indexed inductive sets to justify the interpretation of natural numbers in the scoping model in Section 3. They are also needed for the interpretation of

higher inductive types in Section 5.2 and identity types in Section 5.1. In practise, we will usually not bother to bring a needed fibrant indexed inductive set into the above form and instead work explicitly with the more usual specification in terms of a list of constructors, each taking a certain number non-recursive and recursive arguments. Note that the latter is really an instance of the former since our dependent sums, dependent products, and finite coproducts are extensional (satisfy universal properties). Conversely, the former is an instance of the latter with a single constructor taking a non-recursive and a recursive argument.

C Variations

C.1 Univalence as an axiom

Our treatment extends to the case where the glue types in a cubical cwf as in Section 1.3 are replaced by an operation $\text{Elem}(\Gamma, \text{iUnivalence}_n)$ for $\Gamma : \text{Con}$ and $n \geq 0$, with iUnivalence_n defined in Remark 1.

To define this operation in the scoping model of Section 3, one first shows analogously to Lemmas 5 and 6 that $|-|$ preserves contractible types and that $(A, A') : \text{Type}^*(\Gamma, \Gamma')$ is contractible exactly if A is contractible and $A' \rho \rho' a$ for $\rho : |\Gamma|$ and $\rho' : \Gamma' \rho$ where $a : |A|$ is the induced center of contraction. We have analogous statements for types of homotopy level $n \geq 0$ in \mathcal{M} , in which case we instead have to quantify over all $a : |A|$.

Given $(A, A') : \text{Type}_n^*(\Gamma, \Gamma')$, we have show that the type $(S, S') = \Sigma^*(\mathbb{U}_n, \text{Equiv}^*(\mathbf{q}, A))$ over (Γ, Γ') is contractible in \mathcal{M}^* . Without loss of generality, we may assume the center of contraction of univalence in \mathcal{M} is given by the identity equivalence. Using the observations of the preceding paragraph, it suffices to show that $V' = S' \rho \rho' (\text{pair}(A\rho, (\text{lam}(\mathbf{q}), w)))$ is contractible for $\rho : |\Gamma|$ and $\rho' : \Gamma' \rho$ where w denotes the canonical witness that the identity map $\text{lam}(\mathbf{q})$ on $A\rho$ is an equivalence in \mathcal{M} . Inhabitation is evident, and so it remains to show propositionality. By the case of the preceding paragraph for propositions, the second component of V' is a proposition, and thus we can ignore it for the current goal, which then becomes

$$\text{isProp}(\Sigma(T' : |A| \rightarrow \mathbb{U}_n). \Pi(a : |A|). \text{Equiv}(T' a, A' \rho \rho' a))$$

and follows from univalence in the standard model, justified by glueing.

C.2 Canonicity

A similar scoping argument may be used to provide a reduction-free canonicity argument for cubical type theory with computation rules for filling at type formers alternative to the one of [13]. The key difference is that we now want the filling operation itself to be computable. In the scoping, we then define a type $(A, \langle A', \text{fib}_{A'} \rangle) : \text{Type}(\Gamma, \Gamma')$ to consist of $A : \text{Type}(\Gamma)$ and $A' \rho \rho' a : \mathbb{U}_\omega$ as before, but with $\text{fib}_{A'} \rho \rho' : \mathbb{C}(\Sigma(a : |A| \rho). A' \rho \rho' a)$ such that the first projection relates $\text{fib}_{A'} \rho \rho'$ with the proof of fibrancy of $|A| \rho$. This is easiest formulated by defining an appropriate dependent version $C' : \Pi(A : \mathbb{U}_\omega)(A' : A \rightarrow \mathbb{U}_\omega). \mathbb{C}(A) \rightarrow \mathbb{U}_\omega$ of \mathbb{C} using the methods of Section 2.2.

D Simplicial set model

Choosing for \mathcal{C} the simplex category Δ , for \mathbb{I} the usual interval Δ^1 in simplicial sets, and for \mathbb{F} a small copy $\Omega_{0, \text{dec}}$ of the sublattice of Ω_0 of decidable sieves, we obtain a notion of cubical cwf with a simplicial notion of shape.

Assume now the law of excluded middle. The above choice of $\mathcal{C}, \mathbb{I}, \mathbb{F}$ satisfies all of the assumptions of Section 2.2 but one: the existence of a right adjoint to exponentiation with \mathbb{I} . However, the only place our development makes use of this assumption is in establishing (1). We will instead give a different definition of \mathbb{C} that still satisfies (1). Then the rest of our development still applies to simplicial sets.

A *Kan fibration structure* on a family $Y : X \rightarrow \mathbf{U}_\omega$ in simplicial sets consists of a choice of diagonal fillers in all commuting squares of the form

$$\begin{array}{ccc} \Lambda_k^m & \longrightarrow & \Sigma(x : X).Y x \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta^m & \longrightarrow & X \end{array}$$

with left map a *horn inclusion*. Note that the codomains of horn inclusions are representable. It follows that the presheaf over the slice of simplicial sets over \mathbf{U}_ω is representable. Given $[n] \in \Delta$ and A a presheaf over $\Delta/[n]$, we now define $\mathbb{C}([n], A)$ to be the set of Kan fibration structures on the map $A|_\Delta \rightarrow \Delta^n$. This defines a level preserving map $\mathbb{C} : \mathbf{U}_\omega \rightarrow \mathbf{U}_\omega$. Then the representing object is given by the first projection $\mathbf{U}_\omega^{\text{fib}} \rightarrow \mathbf{U}_\omega$.

Let us now verify (1). Given a simplicial set X with $Y : X \rightarrow \mathbf{U}_\omega$, a global element of $\text{Fill}(X, Y)$ corresponds to a *uniform Kan fibration* structure on $\Sigma(x : X).Y x \rightarrow X$ in the sense of [10]. A uniform Kan fibration structure induces a Kan fibration structure naturally in X , giving the forward direction of (1). For the reverse direction, it suffices to give a uniform Kan fibration structure in the generic case, i.e. a global element of $\text{Fill}(\mathbf{U}_\omega^{\text{fib}}, \lambda(A, c).A)$. This is [10, Theorem 8.9, part (ii)] together with the fact proved in [8, Chapter IV] that Kan fibrations lift against pushout products of interval endpoint inclusions with (levelwise decidable) monomorphisms.⁶

Having verified (1), the rest of our development applies just as well to the case of simplicial sets. In particular, we obtain in the standard model \mathcal{S} of Section 2.3 is a version of the simplicial set model [15] of univalent type theory (using Section 5.1 for identity types).⁷ As per Section 5.2, we furthermore obtain higher inductive types in the simplicial set model in a way that avoids (as suggested by Andrew Swan [25]) the pitfall of fibrant replacement failing to preserve size encountered in [19].

Seeing simplicial sets as a full subtopos of distributive lattice cubical sets as observed in [16], there is a functor from cubical cwfs with $(\mathcal{C}, \mathbb{I}, \mathbb{F}) = (\Delta, \Delta^1, \Omega_{0, \text{dec}})$ to cubical cwfs where \mathcal{C} is the Lawvere theory of distributive lattices, \mathbb{I} is represented by the generic object, and \mathbb{F} is the (small) sublattice of Ω_0 generated by distributive lattice equations. The cubical cwfs in the image of this functor satisfy a *sheaf condition*, which can be represented syntactically as allowing one to e.g. uniquely glue together to a type $\Gamma \vdash_{\{i,j\}} A$ coherent families of types $\Gamma f \vdash_X A_f$ for f a map to X from the free distributive lattice on symbols $\{i, j\}$ such that $f i \leq f j$ or $f j \leq f i$ (compare also the tope logic of [21]).

Applying this functor to the simplicial set model \mathcal{S} discussed above, we obtain an interpretation of distributive lattice cubical type theory (with \mathbb{I} and \mathbb{F} as above) in the sense of the current paper (crucially, without computation rules for filling at type formers) in

⁶ This is the only place where excluded middle is used, to produce a cellular decomposition in terms of simplex boundary inclusions of such a monomorphism.

⁷ Instead of Kan fibration structures, we can work with the property of being a Kan fibration. Then \mathbb{C} is valued in propositions and we would obtain in \mathcal{S} a version of the simplicial set model in which being a type is truly just a property. However, choice would be needed to obtain (1).

23:22 Homotopy canonicity for cubical type theory

homotopy types. Thus, this cubical type theory is homotopically sound in that we can only derive statements which hold in the interpretation in standard homotopy types.